

An elliptic semilinear equation with source term involving boundary measures: the subcritical case.

Marie-Francoise BIDAUT-VERON*

Laurent VIVIER†

Abstract

We study the boundary behaviour of the nonnegative solutions of the semilinear elliptic equation in a bounded regular domain Ω of \mathbb{R}^N ($N \geq 2$),

$$\begin{cases} \Delta u + u^q = 0 & \text{in } \Omega, \\ u = \mu & \text{on } \partial\Omega, \end{cases}$$

where $1 < q < (N+1)/(N-1)$ and μ is a Radon measure on $\partial\Omega$. We give a priori estimates and existence results. They lie on the study of the superharmonic functions in some weighted Marcinkiewicz spaces.

Contents

1	Introduction	2
2	Regularity of the weak solutions	5
2.1	About the Green and Poisson kernels	5
2.2	Regularity of $G(\varphi)$ and $P(\mu)$	8
2.3	Application to the problems 1.1 and 1.4	18
3	Estimate of $G(P^q(\mu))$	19
4	A priori estimates	22
5	Existence results	25

*Laboratoire de Mathématiques et Physique Théorique, CNRS UPRES-A 6083, Faculté des Sciences, Parc de Grandmont, 37200 Tours, France

†Laboratoire de Mathématiques et Physique Théorique, CNRS UPRES-A 6083, Faculté des Sciences, Parc de Grandmont, 37200 Tours, France

1 Introduction

Let Ω be a bounded domain in \mathbb{R}^N ($N \geq 2$) with a C^2 boundary $\partial\Omega$. Here we study the behaviour near the boundary of the nonnegative solutions of the semilinear elliptic equation

$$-\Delta u = u^q \quad \text{in } \Omega, \quad (1.1)$$

where $1 < q < (N+1)/(N-1)$. By solution of (1.1) we mean any function u such that $u^q \in L^1_{loc}(\Omega)$ and satisfying the equation in $\mathcal{D}'(\Omega)$.

We denote by $\rho(x)$ the distance from any point $x \in \Omega$ to $\partial\Omega$, and by $B(x, r)$ the open ball of center x and radius $r > 0$. Let \mathcal{G} be the Green function of the Laplacian in Ω , defined on the set $\{(x, y) \in \overline{\Omega} \times \overline{\Omega} \mid x \neq y\}$. Let \mathcal{P} be the Poisson kernel defined on $\Omega \times \partial\Omega$ by $\mathcal{P}(x, z) = -\partial\mathcal{G}(x, z)/\partial n$. We call $\mathcal{M}(\Omega)$ and $\mathcal{M}(\partial\Omega)$ the spaces of Radon measures on Ω and $\partial\Omega$, and $\mathcal{M}^+(\Omega)$ and $\mathcal{M}^+(\partial\Omega)$ the cones of nonnegative ones.

Observe that any nonnegative and superharmonic function U in Ω satisfies $U \in L^1_{loc}(\Omega)$. From the Herglotz theorem, there exist some unique $\varphi \in \mathcal{M}^+(\Omega)$ and $\mu \in \mathcal{M}^+(\partial\Omega)$ such that U admits the integral representation

$$U = G(\varphi) + P(\mu), \quad (1.2)$$

where, for almost any $x \in \Omega$,

$$G(\varphi)(x) = \int_{\Omega} \mathcal{G}(x, y) d\varphi(y), \quad P(\mu)(x) = \int_{\partial\Omega} \mathcal{P}(x, z) d\mu(z); \quad (1.3)$$

moreover $\int_{\Omega} \rho d\varphi < +\infty$. Reciprocally, for any $\varphi \in \mathcal{M}(\Omega)$ such that $\int_{\Omega} \rho d|\varphi| < +\infty$ and $\mu \in \mathcal{M}(\partial\Omega)$, the function U defined by (1.2) lies in $L^1_{loc}(\Omega)$, and satisfies

$$-\Delta U = \varphi \quad \text{in } \mathcal{D}'(\Omega),$$

see for example [12]. We shall say that U is the *integral solution* of problem

$$\begin{cases} -\Delta U = \varphi & \text{in } \Omega, \\ U = \mu & \text{on } \partial\Omega. \end{cases} \quad (1.4)$$

Hence any solution u of (1.1) in Ω satisfies $\int_{\Omega} \rho u^q dx < +\infty$, and there exists a measure $\mu \in \mathcal{M}^+(\partial\Omega)$ such that

$$\begin{cases} -\Delta u = u^q & \text{in } \Omega, \\ u = \mu & \text{on } \partial\Omega, \end{cases} \quad (1.5)$$

in the integral sense. Our aim is to give *a priori estimates* for any solution of equation (1.1) near the boundary, and also to obtain *existence results* for a given measure μ on $\partial\Omega$.

The problem with the other sign

$$\begin{cases} \Delta u = u^q & \text{in } \Omega, \\ u = \mu & \text{on } \partial\Omega, \end{cases} \quad (1.6)$$

has been studied in [20] and [24] in the subcritical case $1 < q < (N+1)/(N-1)$, and in the supercritical case in [25]. Another approach coming from the probabilistic point of view is done in [14], [15], [22], which gives results in agreement with the previous ones in the case $1 < q \leq 2$. It seems that the probabilistic techniques do not apply to our case. Our approach has to be compared to the methods of P.L. Lions used in [23] for the problem of an interior isolated singularity. Our proofs lie essentially on the study of the superharmonic functions in some weighted Marcinkiewicz spaces.

Let us recall some classical results for the interior problem for a better understanding. Let $x_0 \in \Omega$ and consider any nonnegative solution $w \in C^2(\Omega \setminus \{x_0\})$ of the equation

$$-\Delta w = w^q \quad \text{in } \Omega \setminus \{x_0\}. \quad (1.7)$$

When $1 < q < N/(N-2)$, one can give upper and lower bounds by using Serrin's methods of [28], see for example [4], Lemma A.4. The precise behaviour of w was obtained in [23]. First $w^q \in L^1_{loc}(\Omega)$, and there exists some $\gamma \geq 0$ such that

$$-\Delta w = w^q + \gamma \delta_0, \quad \text{in } \mathcal{D}'(\Omega), \quad (1.8)$$

from the Brézis-Lions Lemma [9]. Then the following estimates hold near x_0 :

$$\gamma E(x_0, x) \leq w(x) \leq \gamma E(x_0, x) (1 + o(1)) \quad (1.9)$$

when $\gamma > 0$, where E is the fundamental solution of the Laplace equation. And the remaining term can be precised according to the values of N, q , see [23]. The function w can be extended as a function $w \in C^2(\Omega)$ if $\gamma = 0$. Concerning the existence of solutions of (1.8) for a given γ , there exists some finite positive γ^* such that the equation (1.8) admits a solution $w \geq 0$, with $w = 0$ on $\partial\Omega$, if and only if $\gamma \in [0, \gamma^*]$. If $q \geq N/(N-2)$, then $\gamma = 0$, see again [23]. If moreover $q \leq (N+2)/(N-2)$, we have the estimate near x_0

$$w(x) \leq C |x - x_0|^{-2/(q-1)} \quad (1.10)$$

with $C = C(N, q)$, see [17] and [11].

Now let us come back to the boundary problem. As in [20] we can define another concept of solution. Let $C^{1,1}_0(\bar{\Omega})$ be the space of C^1 functions vanishing on $\partial\Omega$ with Lipschitz continuous gradient. For any $\varphi \in \mathcal{M}(\Omega)$ such that $\int_{\Omega} \rho d|\varphi| < +\infty$ and any $\mu \in \mathcal{M}(\partial\Omega)$, we shall say that a function U is *weak solution* of problem (1.4) if $U \in L^1(\Omega)$ and

$$\int_{\Omega} U(-\Delta \xi) dx = \int_{\Omega} \xi d\varphi - \int_{\partial\Omega} \frac{\partial \xi}{\partial n} d\mu \quad (1.11)$$

for any $\xi \in C_0^{1,1}(\overline{\Omega})$. In **Section 2**, we first verify that the integral solution coincides with the weak one, and hence is in $L^1(\Omega)$. Then we give regularity results of the general weak solution U of (1.4) in some Marcinkiewicz spaces with a weight of the form ρ^β ($\beta \in \mathbb{R}$). They lie on precise estimates of the Green and Poisson kernel. Up to our knowledge, most of them are new, more especially as the measure φ may be unbounded, and can present an interest in themselves. They are fundamental to obtain a priori estimates and existence results for the problem (1.5), above all in the most delicate case $N/(N-1) \leq q < (N+1)/(N-1)$.

In **Section 3**, we give an a priori estimate for the function $G(P^q(\mu))$, for any $\mu \in \mathcal{M}^+(\partial\Omega)$:

Theorem 1.1 *Assume that $1 < q < (N+1)/(N-1)$. Then $P^q(\mu) \in L^1(\Omega, \rho dx)$ for any $\mu \in \mathcal{M}^+(\partial\Omega)$, and there exists a constant $K = K(N, \Omega, q)$ such that,*

$$G(P^q(\mu)) \leq K \mu(\partial\Omega)^{q-1} P(\mu) \quad \text{in } \Omega. \quad (1.12)$$

This result is interesting from two points of view. Above all it allows to construct supersolutions, hence to get existence results. Concerning the a priori estimates for (1.5), setting $u = P(\mu) + v$, the function v satisfies

$$v = G(u^q) \leq 2^{q-1} [G(P^q(\mu)) + G(v^q)], \quad \text{a.e. in } \Omega,$$

hence any estimate on $G(P^q(\mu))$ gives informations on v .

In **Section 4** we prove our main result, which is an a priori estimate of any solution of (1.5) in terms of the solution $P(\mu)$ of the associated linear problem. It lies on the results of Section 2. It also uses the estimate (1.12), which in fact can be shown almost as a necessary condition of existence of solutions, by using recent techniques of ([8]).

Theorem 1.1 *Assume that $1 < q < (N+1)/(N-1)$. Let $\mu \in \mathcal{M}^+(\partial\Omega)$, and u be any nonnegative solution of (1.5). Then there exists a constant $C = C(N, q, \Omega, \mu(\partial\Omega))$, such that*

$$P(\mu) \leq u \leq C (P(\mu) + \rho) \quad \text{in } \Omega \quad (1.13)$$

(and $u \in C^\infty(\Omega) \cap C^{1,\alpha}(\overline{\Omega})$ for any $\alpha \in (0, 1)$ if $\mu = 0$).

Theorem 1.2 *More precisely, if $\mu = \sigma \delta_a$ for some $a \in \partial\Omega$ and $\sigma > 0$, then near the point a ,*

$$\sigma P(\delta_a)(x) \leq u(x) \leq \sigma P(\delta_a)(x) \left[1 + O(|x - a|^{N+1-(N-1)q}) \right]. \quad (1.14)$$

This result applies in particular to any solution u of (1.1), such that $u \in C(\overline{\Omega} \setminus \{a\})$ and $u = 0$ on $\partial\Omega \setminus \{a\}$, since its trace is necessarily of the form $\mu = \sigma \delta_a$ for some $\sigma \geq 0$. Notice also that in case $q < (N+1)/(N-1)$, Theorem 1.1 extends some a priori estimates of [13], [18] to the case of unbounded boundary data.

In **Section 5**, we use for proving our second main theorem, which gives existence results.

Theorem 1.2 *Assume that $1 < q < (N+1)/(N-1)$. Let $\mu \in \mathcal{M}^+(\partial\Omega)$ with $\mu(\partial\Omega) = 1$, and $\sigma \geq 0$. Then there exists some finite positive σ^* such that the problem*

$$\mathcal{S}_\sigma \quad \begin{cases} -\Delta u = u^q & \text{in } \Omega, \\ u = \sigma \mu & \text{on } \partial\Omega, \end{cases} \quad (1.15)$$

admits a solution if and only if $\sigma \in [0, \sigma^]$.*

The existence of solutions for small σ is a direct consequence of Theorem 1.1. The existence of an interval $[0, \sigma^*]$ is an adaptation of some results of ([8]).

In conclusion, in the subcritical case we have completely extended the results of an interior punctual singularity to any boundary measure singularity. The next step, that is the study of the case $q \geq (N+1)/(N-1)$, is still open.

Note added in proof. In the moment this article was in printing, we received a preprint of H. Amann and P. Quittner [1], where they consider more general problems with interior and boundary bounded measures, and use duality methods. In case of problem (1.5), they get a regularity result in $W^{1-\varepsilon,1}(\Omega)$ for any $\varepsilon \in (0, 1)$, and prove the existence of at least two solutions, under the condition $q < N/(N-1)$.

2 Regularity of the weak solutions

2.1 About the Green and Poisson kernels

Here we recall and complete some classical estimates for the Green function and the Poisson kernel. For almost any $y \in \Omega$ and $z \in \partial\Omega$, the functions $\mathcal{G}(\cdot, y)$ and $\mathcal{P}(\cdot, z)$ are the integral solutions of

$$\begin{cases} -\Delta \mathcal{G}(\cdot, y) = \delta_y & \text{in } \Omega, \\ \mathcal{G}(\cdot, y) = 0 & \text{on } \partial\Omega, \end{cases} \quad \begin{cases} -\Delta \mathcal{P}(\cdot, z) = 0 & \text{in } \Omega, \\ \mathcal{P}(\cdot, z) = \delta_z & \text{on } \partial\Omega, \end{cases}$$

where δ_y, δ_z are the Dirac masses at points $y \in \Omega$, and $z \in \partial\Omega$.

Proposition 2.1 *There exists a constant $c_N = c_N(N, \Omega)$ such that*

i) for any $(x, y) \in \Omega \times \Omega$ with $x \neq y$,

$$\mathcal{G}(x, y) \leq \begin{cases} c_N |x - y|^{2-N} & \text{if } N \geq 3, \\ c_2 (1 + |\ln |x - y||) & \text{if } N = 2, \end{cases} \quad (2.1)$$

$$\mathcal{G}(x, y) \leq c_N \rho(x) |x - y|^{1-N}, \quad (2.2)$$

$$\mathcal{G}(x, y) \leq c_N \rho(x) \rho(y) |x - y|^{-N}, \quad (2.3)$$

$$\mathcal{G}(x, y) \leq \begin{cases} c_N (\rho(x)/\rho(y)) |x - y|^{2-N} & \text{if } N \geq 3, \\ c_2 (\rho(x)/\rho(y))(1 + |\ln |x - y||) & \text{if } N = 2, \end{cases} \quad (2.4)$$

and

$$|\nabla_x \mathcal{G}(x, y)| \leq c_N |x - y|^{1-N}, \quad (2.5)$$

$$|\nabla_x \mathcal{G}(x, y)| \leq c_N \rho(y) |x - y|^{-N}, \quad (2.6)$$

$$|\nabla_x \mathcal{G}(x, y)| \leq c_N (\rho(y)/\rho(x)) |x - y|^{1-N}. \quad (2.7)$$

ii) For any $(x, z) \in \Omega \times \partial\Omega$,

$$c_N^{-1} \rho(x) |x - z|^{-N} \leq \mathcal{P}(x, z) \leq c_N \rho(x) |x - z|^{-N} \leq c_N |x - z|^{1-N}, \quad (2.8)$$

$$|\nabla_x \mathcal{P}(x, z)| \leq c_N |x - z|^{-N}. \quad (2.9)$$

Proof. But for (2.4) and (2.7), all these estimates are well known. They are deduced from the explicit expression of \mathcal{G} in an half-space, and extended to any C^2 bounded open set. For the lower estimate of (2.8), see [21]. Let us prove (2.4): it is a consequence of (2.1) and (2.2). Indeed that is true in the set $\{\rho(y) \leq 2\rho(x)\}$. Now suppose $\rho(y) > 2\rho(x)$. Let $x^* \in \partial\Omega$ such that $|x - x^*| = \rho(x)$. Then $|x - y| \geq |x^* - y| - |x^* - x| \geq \rho(y) - \rho(x) \geq \rho(y)/2$, hence (2.2) implies if $N \geq 3$

$$\mathcal{G}(x, y) \leq c_N \frac{\rho(x)}{|x - y|} |x - y|^{2-N} \leq 2c_N \frac{\rho(x)}{\rho(y)} |x - y|^{2-N};$$

thus (2.4) holds with a new constant c_N ; likewise if $N = 2$. Similarly (2.5) and (2.6) imply (2.7). ■

Remark 2.1 Notice that (2.2) can be deduced from (2.1) and (2.5) when $N \geq 3$. Indeed (2.1) implies (2.2) in the set $\{|x - y| \leq 2\rho(x)\}$. Now suppose that $|x - y| > 2\rho(x)$. Defining x^* as above, we have $[x, x^*) \subset \Omega$, and from (2.5),

$$\begin{aligned} \mathcal{G}(x, y) &= |\mathcal{G}(x, y) - \mathcal{G}(x^*, y)| \leq |x - x^*| \sup_{t \in [x, x^*)} |\nabla_x \mathcal{G}(t, y)| \\ &\leq c_N \rho(x) \sup_{t \in [x, x^*)} |y - t|^{1-N} \leq 2^{N-1} c_N \rho(x) |x - y|^{1-N}, \end{aligned}$$

since $|y - t| \geq |y - x| - |t - x| \geq |y - x| - \rho(x) \geq |x - y|/2$. Hence (2.2) holds. Similarly (2.2) and (2.6) imply (2.3) for any $N \geq 2$. And (2.6) also implies the

upper estimate (2.8), since $\mathcal{P}(x, z) = -\partial\mathcal{G}(x, y)/\partial n$ and \mathcal{G} is of class C^1 in the set $\{(x, y) \in \overline{\Omega} \times \overline{\Omega} \mid x \neq y\}$. The estimates (2.2), (2.5) and (2.6) are proved in [33] in the more general framework of a Lyapounov open set. And (2.9) is proved in a $C^{1,\alpha}$ open set in [26].

As a consequence we can compare the integral and weak solutions of (1.4):

Corollary 2.2 *For any $\varphi \in \mathcal{M}(\Omega)$ such that $\int_{\Omega} \rho \, d|\varphi| < +\infty$ and any $\mu \in \mathcal{M}(\partial\Omega)$, a function U is weak solution of problem (1.4) if and only if it is given by the representation (1.2). Consequently, (1.4) has a unique weak solution U in $L^1(\Omega)$, and*

$$\|U\|_{L^1(\Omega)} \leq C \left(\int_{\Omega} \rho \, d|\varphi| + \int_{\partial\Omega} d|\mu| \right), \quad (2.10)$$

for some constant $C = C(N, \Omega)$.

Proof. By considering the positive and negative parts of φ and μ , we can assume that the two measures are nonnegative. Let us prove that the integral solution U is a weak solution. The main point is to prove that $U \in L^1(\Omega)$. From (2.2) and (2.8),

$$\int_{\Omega \times \Omega} \mathcal{G}(x, y) \, dx \, d\varphi(y) \leq c_N \int_{\Omega} \left(\int_{\Omega} |x - y|^{1-N} \, dx \right) \rho(y) \, d\varphi(y) \leq C \int_{\Omega} \rho \, d\varphi, \quad (2.11)$$

with another constant $C = C(N, \Omega)$, and

$$\int_{\Omega \times \partial\Omega} \mathcal{P}(x, z) \, dx \, d\mu(z) \leq c_N \int_{\Omega} \left(\int_{\partial\Omega} |x - z|^{1-N} \, dx \right) d\mu(z) \leq C \int_{\partial\Omega} d\mu, \quad (2.12)$$

hence $U \in L^1(\Omega)$ and

$$\int_{\Omega} U(x) \, dx \leq C \left(\int_{\Omega} \rho \, d\varphi + \int_{\partial\Omega} d\mu \right).$$

Now for any $\xi \in C_0^{1,1}(\overline{\Omega})$, we have

$$\mathcal{G}(x, y) \, \Delta\xi(x) \in L^1(\Omega \times \Omega, dx \, d\varphi(y)), \quad \mathcal{P}(x, z) \, \Delta\xi(x) \in L^1(\Omega \times \partial\Omega, dx \, d\mu(z))$$

from (2.11) and (2.12). Then U is a weak solution from the Fubini theorem, and (2.10) follows. Reciprocally, if U is a weak solution of problem (1.4), then $-\Delta U = \varphi$ in $\mathcal{D}'(\Omega)$, and there exists a unique measure $\tilde{\mu} \in \mathcal{M}^+(\partial\Omega)$ such that $U = G(\varphi) + P(\tilde{\mu})$. Then U is a weak solution for the problem with data φ and $\tilde{\mu}$. Hence for any $\xi \in C_0^{1,1}(\overline{\Omega})$,

$$\int_{\partial\Omega} \frac{\partial\xi}{\partial n} \, d\mu = \int_{\partial\Omega} \frac{\partial\xi}{\partial n} \, d\tilde{\mu},$$

which implies $\tilde{\mu} = \mu$. Then U is the integral solution of (1.4), and (2.10) follows again. ■

Remark 2.2 Thus for any $\varphi \in \mathcal{M}(\Omega)$ such that $\int_{\Omega} \rho d|\varphi| < +\infty$ and any $\mu \in \mathcal{M}(\partial\Omega)$, the problem (1.4) is well posed in $L^1(\Omega)$. We find again in a very short way the result of [7] where φ is a measurable function with $\rho \varphi \in L^1(\Omega)$ and $\mu \in L^1(\partial\Omega)$.

The lower estimate of the Poisson kernel (2.8) also shows that the value $q = (N+1)/(N-1)$ is a natural barrier for the problem (1.5):

Corollary 2.3 *Assume that $q \geq (N+1)/(N-1)$. Then problem (1.5) has no solution for a positive measure concentrated at some point $a \in \partial\Omega$.*

Proof. For any $\mu \in \mathcal{M}^+(\partial\Omega)$, if (1.5) has a solution u , then $u \geq P(\mu)$, but $u^q \in L^1(\Omega, \rho dx)$, hence also $P^q(\mu)$. Suppose that $\mu > 0$ with $\text{supp } \mu = \{a\}$, that means $\mu = \sigma \delta_a$ for some $\sigma > 0$. From (2.8), we have

$$\begin{aligned} \int_{\Omega} P^q(\delta_a) \rho dx &\geq c_N^{-q} \int_{\Omega} |x-a|^{-Nq} \rho^{q+1} dx \\ &\geq 2^{-(q+1)} c_N^{-q} \int_{\{x \in \Omega \mid \rho(x) \geq |x-a|/2\}} |x-a|^{q+1-Nq} dx \end{aligned}$$

But the set $\{x \in \Omega \mid \rho(x) \geq |x-a|/2\}$ contains the intersection of a cone of vertex a and angle $\pi/3$ with a small ball of center a . Hence the integral is divergent, since $q \geq (N+1)/(N-1)$. Then we arrive to a contradiction. ■

2.2 Regularity of $G(\varphi)$ and $P(\mu)$

Now we are going to complete the estimate (2.10) by much more precise estimates of the functions $G(\varphi)$ and $P(\mu)$ in Marcinkiewicz weighted spaces, with a power of the distance ρ as a weight function. Let us recall their definition. For any $k \in \mathbb{R}$ with $k \geq 1$, and any positive weight function $\eta \in C(\Omega)$, we denote by $L^k(\Omega, \eta dx)$ the space of measurable functions v on Ω such that

$$\|v\|_{L^k(\Omega, \eta dx)} = \left(\int_{\Omega} |v|^k \eta dx \right)^{1/k} < +\infty,$$

and the Marcinkiewicz space $M^k(\Omega, \eta dx)$ is the space of measurable functions v on Ω such that

$$\sup_{\lambda > 0} \lambda \left(\int_{\{x \in \Omega \mid |v(x)| \geq \lambda\}} \eta dx \right)^{1/k} < +\infty.$$

And for any $k > 1$, $M^k(\Omega, \eta dx)$ is also the normed space of the v such that

$$\|v\|_{M^k(\Omega, \eta dx)} = \sup \left[\left(\int_{\omega} |v| \eta dx \right) / \left(\int_{\omega} \eta dx \right)^{1-1/k} \right] < +\infty,$$

where the supremum is taken over the measurable subsets ω of Ω such that $\int_{\omega} \eta \, dx$ is finite. We have $L^k(\Omega, \eta \, dx) \subset M^k(\Omega, \eta \, dx)$. If $\eta \in L^1(\Omega)$ (in particular $\eta = \rho^\beta$ with $\beta > -1$), then

$$M^k(\Omega, \eta \, dx) \subset L^m(\Omega, \eta \, dx) \quad \text{for any } m \in [1, k].$$

If $\eta \equiv 1$, $L^k(\Omega, \eta \, dx) = L^k(\Omega)$, and $M^k(\Omega, \eta \, dx) = M^k(\Omega)$.

Recall that the solution w of the interior problem (1.7) lies in $M_{loc}^{N/(N-2)}(\Omega)$ if $N \geq 3$, and in $L_{loc}^p(\Omega)$ for any $p \geq 1$ if $N = 2$, see [9]. On the other part, from [20] and Corollary 2.2, for any nonnegative $\mu \in L^1(\partial\Omega)$, the function $P(\mu)$ lies in $M^{N/(N-1)}(\Omega) \cap M^{(N+1)/(N-1)}(\Omega, \rho \, dx)$. The following Lemma extends the techniques used in [2] and [20]:

Lemma 2.4 *Let ν be a nonnegative bounded Radon measure on $D = \Omega$ or $\partial\Omega$, and $\eta \in C(\Omega)$ be a positive weight function. Let \mathcal{H} be a continuous nonnegative function on $\{(x, t) \in \Omega \times D \mid x \neq t\}$. For any $\lambda > 0$, we set*

$$A_\lambda(t) = \{x \in \Omega \setminus \{t\} \mid \mathcal{H}(x, t) > \lambda\}, \quad (2.13)$$

$$m_\lambda(t) = \int_{A_\lambda(t)} \eta \, dx. \quad (2.14)$$

Suppose that for some $C \geq 0$ and $k > 1$

$$m_\lambda(t) \leq C \lambda^{-k}, \quad \forall \lambda > 0. \quad (2.15)$$

Then the function

$$x \in \Omega \mapsto H(x) = \int_D \mathcal{H}(x, t) \, d\nu(t) \quad (2.16)$$

is in $M^k(\Omega, \eta \, dx)$ and

$$\|H\|_{M^k(\Omega, \eta \, dx)} \leq (1 + \frac{k}{k-1}C) \nu(D). \quad (2.17)$$

Proof. Let ω be any measurable subset of Ω such that $\int_{\omega} \eta \, dx$ is finite. Then for any $\lambda > 0$, and any $t \in D$,

$$\int_{\omega} \mathcal{H}(x, t) \eta(x) \, dx \leq \int_{A_\lambda(t)} \mathcal{H}(x, t) \eta(x) \, dx + \lambda \int_{\omega} \eta(x) \, dx,$$

with

$$\begin{aligned} \int_{A_\lambda(t)} \mathcal{H}(x, t) \eta(x) \, dx &= - \int_{\lambda}^{+\infty} \theta \, dm_\theta(t) = \lambda m_\lambda(t) + \int_{\lambda}^{+\infty} m_\theta(t) \, d\theta \\ &\leq \frac{k}{k-1} C \lambda^{1-k}. \end{aligned}$$

Choosing $\lambda = \left(\int_{\omega} \eta \, dx \right)^{-1/k}$, we get

$$\int_{\omega} \mathcal{H}(x, t) \eta(x) \, dx \leq \left(1 + \frac{k}{k-1} C \right) \left(\int_{\omega} \eta \, dx \right)^{1-1/k},$$

and by integration over D with respect to the measure ν ,

$$\begin{aligned} \int_{\omega} H(x) \eta(x) \, dx &= \int_{\omega} \int_D \mathcal{H}(x, t) \eta(x) \, dx \, d\nu(t) \\ &\leq \left(1 + \frac{k}{k-1} C \right) \nu(D) \left(\int_{\omega} \eta \, dx \right)^{1-1/k}, \end{aligned}$$

hence the conclusion. \blacksquare

Let us first complete the estimates of [20] for the function $P(\mu)$:

Theorem 2.5 *For any $\mu \in \mathcal{M}(\partial\Omega)$, let $\Psi = P(\mu)$ be the solution of the problem*

$$\begin{cases} -\Delta \Psi = 0 & \text{in } \Omega, \\ \Psi = \mu & \text{on } \partial\Omega. \end{cases} \quad (2.18)$$

Then

$$\Psi \in M^{(N+\beta)/(N-1)}(\Omega, \rho^{\beta} \, dx) \quad (2.19)$$

for any $\beta > -1$, and

$$|\nabla \Psi| \in M^{(N+\gamma)/N}(\Omega, \rho^{\gamma} \, dx), \quad (2.20)$$

for any $\gamma > 0$. Moreover there exists constants $C = C(\Omega, N, \beta) > 0$ and $C' = C'(\Omega, N, \gamma) > 0$ such that

$$\|\Psi\|_{M^{(N+\beta)/(N-1)}(\Omega, \rho^{\beta} \, dx)} \leq C \, |\mu|(\partial\Omega), \quad (2.21)$$

$$\| |\nabla \Psi| \|_{M^{(N+\gamma)/N}(\Omega, \rho^{\gamma} \, dx)} \leq C' \, |\mu|(\partial\Omega). \quad (2.22)$$

Proof. First step: estimate of the function. We can suppose that μ is nonnegative. Let β be a real parameter. We shall apply Lemma 2.4 with

$$D = \partial\Omega, \quad \eta = \rho^{\beta}, \quad \nu = \mu, \quad \text{and} \quad \mathcal{H}(x, t) = \mathcal{P}(x, t). \quad (2.23)$$

From (2.8), for any $t \in \partial\Omega$, and any $\lambda > 0$, and any $x \in A_{\lambda}(t)$,

$$\lambda \leq c_N \rho(x) |x - t|^{-N} \leq c_N |x - t|^{1-N}.$$

Hence if $\beta \geq 0$,

$$m_{\lambda}(t) \leq \int_{B(t, (c_N/\lambda)^{1/(N-1)})} \rho^{\beta} \, dx \leq \int_{B(t, (c_N/\lambda)^{1/(N-1)})} |x - t|^{\beta} \, dx \leq C \lambda^{-(N+\beta)/(N-1)}.$$

If $\beta < 0$, then

$$\begin{aligned} m_\lambda(t) &\leq \int_{B(t, (c_N/\lambda)^{1/(N-1)})} \rho^\beta dx \leq \int_{B(t, (c_N/\lambda)^{1/(N-1)})} (\lambda |x - t|^N / c_N)^\beta dx \\ &\leq C \lambda^\beta \int_0^{(c_N/\lambda)^{1/(N-1)}} r^{N-1+N\beta} dr \leq C \lambda^{-(N+\beta)/(N-1)}, \end{aligned}$$

under the condition $\beta > -1$. Then Lemma 2.4 gives (2.19) and (2.21).

Second step: estimate of the gradient. Let $i \in \{1, \dots, N\}$. Here we use Lemma 2.4 with

$$D = \partial\Omega, \quad \eta = \rho^\gamma, \quad \nu = \mu, \quad \text{and} \quad \mathcal{H}(x, t) = \partial\mathcal{P}(x, t)/\partial x_i. \quad (2.24)$$

From (2.9), for any $t \in \partial\Omega$, and any $\lambda > 0$, and any $x \in A_\lambda(t)$,

$$\lambda \leq c_N |x - t|^{-N}.$$

Then if $\gamma \geq 0$,

$$m_\lambda(t) \leq \int_{B(t, (c_N/\lambda)^{1/N})} \rho^\gamma dx \leq \int_{B(t, (c_N/\lambda)^{1/N})} |x - t|^\gamma dx \leq C \lambda^{-(N+\gamma)/N}.$$

Hence if $\gamma > 0$, the function

$$Q_i(x) = \int_{\partial\Omega} \frac{\partial\mathcal{P}(x, t)}{\partial x_i} d\mu(t)$$

lies in $M^{(N+\gamma)/N}(\Omega, \rho^\gamma dx)$. But $Q_i = \partial P(\mu)/\partial x_i$ from the derivation theorem, so that (2.20) and (2.22) hold. ■

Let us now give precise estimates of $G(\varphi)$. They are one of the keys of Theorem 1.1.

Theorem 2.6 *For any $\varphi \in \mathcal{M}(\Omega)$ such that $\int_\Omega \rho^\alpha d|\varphi| < +\infty$, with $\alpha \in [0, 1]$, let $\Phi = G(\varphi)$ be the solution of problem*

$$\begin{cases} -\Delta\Phi = \varphi & \text{in } \Omega, \\ \Phi = 0 & \text{on } \partial\Omega. \end{cases} \quad (2.25)$$

i) Then if $N \geq 3$,

$$\Phi \in M^{(N+\beta)/(N-2+\alpha)}(\Omega, \rho^\beta dx) \quad (2.26)$$

for any $\beta \in (-N/(N+\alpha-1), \alpha N/(N-2))$ if $\alpha \neq 0$, for any $\beta \in (-N/(N-1), 0]$ if $\alpha = 0$. In any case, there exists some $C = C(\Omega, N, \alpha, \beta) > 0$ such that

$$\|\Phi\|_{M^{(N+\beta)/(N-2+\alpha)}(\Omega, \rho^\beta dx)} \leq C \int_\Omega \rho^\alpha d|\varphi|, \quad (2.27)$$

If $N = 2$, and $\alpha \in (0, 1]$,

$$\Phi \in M^{(2+\beta-\varepsilon)/\alpha}(\Omega, \rho^\beta dx) \quad (2.28)$$

for any $\beta \in (-2/(1+\alpha), +\infty)$ and $\varepsilon > 0$ small enough; if $\alpha = 0$, then

$$\Phi \in M^p(\Omega, \rho^\beta dx) \quad (2.29)$$

for any $\beta \in (-2, 0]$ and $p \in (\max(1, -\beta), +\infty)$; with similar continuity properties in those spaces.

ii) For any $N \geq 2$,

$$|\nabla \Phi| \in M^{(N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx), \quad \Phi/\rho \in M^{(N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx), \quad (2.30)$$

for any $\gamma \in [0, \alpha N/(N-1))$ if $\alpha \in (0, 1)$, any $\gamma \in (0, N/(N-1))$ if $\alpha = 1$, and $\gamma = 0$ if $\alpha = 0$, and there exists some $C' = C(\Omega, N, \alpha, \gamma) > 0$ such that

$$\| |\nabla \Phi| + \Phi/\rho \|_{M^{(N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx)} \leq C' \int_{\Omega} \rho^\alpha d|\varphi|. \quad (2.31)$$

Proof. First step: estimate of the function. Here also we can assume that φ is nonnegative. Let $\alpha \in [0, 1]$ be fixed, and β be a real parameter. We have

$$G(\varphi)(x) = \int_{\Omega} \frac{\mathcal{G}(x, y)}{\rho^\alpha(y)} \rho^\alpha(y) d\varphi(y),$$

We shall apply Lemma 2.4 with

$$D = \Omega, \quad \eta = \rho^\beta, \quad \nu = \rho^\alpha \varphi, \quad \text{and} \quad \mathcal{H}(x, t) = \mathcal{G}(x, t)/\rho^\alpha(t). \quad (2.32)$$

i) First assume $N \geq 3$. From (2.1) and (2.2), for any $x, t \in \Omega$ with $x \neq t$,

$$\mathcal{G}(x, t) \leq c_N |x - t|^{(2-N)(1-\alpha)} (\rho(t) |x - t|^{1-N})^\alpha \leq c_N \rho^\alpha(t) |x - t|^{2-N-\alpha}, \quad (2.33)$$

Moreover, from (2.1) and (2.4),

$$\mathcal{G}(x, t) \leq c_N |x - t|^{(2-N)(1-\alpha)} \left(\frac{\rho(t)}{\rho(x)} |x - t|^{2-N} \right)^\alpha \leq c_N \frac{\rho^\alpha(t)}{\rho^\alpha(x)} |x - t|^{2-N}, \quad (2.34)$$

and from (2.2) and (2.3),

$$\mathcal{G}(x, t) \leq c_N (\rho(x) |x - t|^{1-N})^{(1-\alpha)} (\rho(x) \rho(t) |x - t|^{-N})^\alpha \leq c_N \rho(x) \rho^\alpha(t) |x - t|^{1-N-\alpha}. \quad (2.35)$$

Then for any $\lambda > 0$, and any $x \in A_\lambda(t)$, from (2.33)

$$\lambda \leq c_N |x - t|^{2-N-\alpha}, \quad (2.36)$$

and from (2.34) and (2.35),

$$\rho^\alpha(x) \leq (c_N/\lambda) |x - t|^{2-N}, \quad \text{and} \quad \rho(x) \geq (\lambda/c_N) |x - t|^{N-1+\alpha}. \quad (2.37)$$

First suppose that $\alpha > 0$ and $\beta > 0$. Then

$$\begin{aligned} m_\lambda(t) &\leq \int_{B(t, (c_N/\lambda)^{1/(N-2+\alpha)})} \rho^\beta dx \leq \int_{B(t, (c_N/\lambda)^{1/(N-2+\alpha)})} ((c_N/\lambda) |x - t|^{2-N})^{\beta/\alpha} dx \\ &\leq C \lambda^{-\beta/\alpha} \int_0^{(c_N/\lambda)^{1/(N-2+\alpha)}} r^{N-1-(N-2)\beta/\alpha} dr \leq C \lambda^{-(N+\beta)/(N-2+\alpha)}, \end{aligned}$$

under the condition $\beta < \alpha N/(N-2)$. Now suppose that $\beta \leq 0$. Then

$$\begin{aligned} m_\lambda(t) &\leq \int_{B(t, (c_N/\lambda)^{1/(N-2+\alpha)})} \rho^\beta dx \leq \int_{B(t, (c_N/\lambda)^{1/(N-2+\alpha)})} ((\lambda/c_N) |x - t|^{N-1+\alpha})^\beta dx \\ &\leq C \lambda^\beta \int_0^{(c_N/\lambda)^{1/(N-2+\alpha)}} r^{N-1+(N-1+\alpha)\beta} dr \leq C \lambda^{-(N+\beta)/(N-2+\alpha)}, \end{aligned}$$

under the condition $\beta > -N/(N-1+\alpha)$. Hence Lemma 2.4 applies and gives the estimates (2.26) and (2.27) for Φ .

ii) Now assume $N = 2$. Then (2.33) and (2.34) are replaced by

$$\mathcal{G}(x, t) \leq c_2 (1 + |\ln |x - t||)^{(1-\alpha)} (\rho(t) |x - t|^{-1})^\alpha \quad (2.38)$$

$$\mathcal{G}(x, t) \leq c_2 \frac{\rho^\alpha(t)}{\rho^\alpha(x)} (1 + |\ln |x - t||), \quad (2.39)$$

and (2.35) is still valid. Then (2.36) and (2.37) become

$$\lambda \leq c_2 |x - t|^{-\alpha} (1 + |\ln |x - t||)^{(1-\alpha)} \quad (2.40)$$

$$\rho^\alpha(x) \leq (c_2/\lambda) (1 + |\ln |x - t||), \quad \text{and} \quad \rho(x) \geq (\lambda/c_2) |x - t|^{1+\alpha}. \quad (2.41)$$

First suppose $\alpha \in (0, 1]$. Notice that (2.40) and (2.41) imply for any $\varepsilon > 0$,

$$\lambda \leq C_\varepsilon |x - t|^{-\alpha-\varepsilon}, \quad \rho^\alpha(x) \leq (C_\varepsilon/\lambda) |x - t|^{-\varepsilon},$$

with $C_\varepsilon = C_\varepsilon(N, \Omega, \varepsilon)$, since Ω is bounded. If $\beta > 0$, then

$$m_\lambda(t) \leq \int_{B(t, (C_\varepsilon/\lambda)^{1/(\alpha+\varepsilon)})} \rho^\beta dx \leq \int_{B(t, (C_\varepsilon/\lambda)^{1/(\alpha+\varepsilon)})} ((C_\varepsilon/\lambda) |x - t|^{-\varepsilon})^{\beta/\alpha} dx$$

so that for any small $\varepsilon > 0$,

$$m_\lambda(t) \leq C'_\varepsilon \lambda^{-(2+\beta-\varepsilon)/\alpha}$$

with $C'_\varepsilon = C'_\varepsilon(\Omega, \alpha, \beta, \varepsilon)$, hence $\Phi \in M^{(2+\beta-\varepsilon)/\alpha}(\Omega, \rho^\beta dx)$. In case $\beta \leq 0$, we find

$$\begin{aligned} m_\lambda(t) &\leq \int_{B(t, (C_\varepsilon/\lambda)^{1/(\alpha+\varepsilon)})} \rho^\beta dx \leq C_\varepsilon \lambda^\beta \int_0^{(C_\varepsilon/\lambda)^{1/(\alpha+\varepsilon)}} r^{1+(1+\alpha)\beta} dr \\ &\leq C'_\varepsilon \lambda^{\beta-(2+(1+\alpha)\beta)/(\alpha+\varepsilon)}, \end{aligned}$$

with $C''_\varepsilon = C''_\varepsilon(\Omega, \alpha, \beta, \varepsilon)$, under the condition $\beta > -2/(1+\alpha)$, hence the same conclusion. Now suppose $\alpha = 0$ and $-2 < \beta \leq 0$. Observe that the condition (2.40) implies $|x-t| \leq C_\Omega e^{-\lambda/c_2}$, with for example $C_\Omega = e(1 + (\text{diam } \Omega)^2)$. Then from (2.41),

$$m_\lambda(t) \leq \int_{B(t, C_\Omega e^{-\lambda/c_2})} \rho^\beta dx \leq C \lambda^\beta \int_0^{C_\Omega e^{-\lambda/c_2}} r^{1+\beta} dr \leq C \lambda^\beta e^{-(\beta+2)\lambda/c_2} \leq C_p \lambda^{-p}$$

for any $p > -\beta$, hence $\Phi \in M^p(\Omega, \rho^\beta dx)$ for any $p \in (\max(1, -\beta), +\infty)$.

Second step: estimate of the gradient and of Φ/ρ . In the same way, we take

$$\begin{aligned} D &= \Omega, \quad \eta = \rho^\gamma, \quad \nu = \rho^\alpha \varphi, \quad \text{and} \\ \mathcal{H}(x, t) &= \frac{\partial \mathcal{G}(x, t)}{\partial x_i} / \rho^\alpha(t) \quad \text{or} \quad \mathcal{H}(x, t) = \mathcal{G}(x, t) / \rho(x) \rho^\alpha(t) \end{aligned} \quad (2.42)$$

As above, for any $N \geq 2$, from (2.5) and (2.6), for any $x, t \in \Omega$ with $x \neq t$,

$$\left| \frac{\partial \mathcal{G}(x, t)}{\partial x_i} \right| \leq c_N \rho^\alpha(t) |x-t|^{1-N-\alpha}, \quad (2.43)$$

and similarly from (2.2) and (2.3),

$$\mathcal{G}(x, t) / \rho(x) \leq c_N \rho^\alpha(t) |x-t|^{1-N-\alpha}. \quad (2.44)$$

And from (2.5) and (2.7)

$$\left| \frac{\partial \mathcal{G}(x, t)}{\partial x_i} \right| \leq c_N \frac{\rho^\alpha(t)}{\rho^\alpha(x)} |x-t|^{1-N}, \quad (2.45)$$

and similarly from (2.2), which is symmetrical in x and y ,

$$\mathcal{G}(x, t) / \rho(x) \leq \frac{\rho^\alpha(t)}{\rho^\alpha(x)} |x-t|^{1-N}. \quad (2.46)$$

Then for any $\lambda > 0$, and any $x \in A_\lambda(t)$, from (2.43) and (2.45), or from (2.44) and (2.46),

$$\lambda \leq c_N |x-t|^{1-N-\alpha}, \quad \text{and} \quad \rho^\alpha(x) \leq (c_N/\lambda) |x-t|^{1-N},$$

First assume that $\alpha > 0$ and $\gamma \geq 0$. Then

$$\begin{aligned} m_\lambda(t) &\leq \int_{B(t, (c_N/\lambda)^{1/(N-1+\alpha)})} \rho^\gamma dx \leq \int_{B(t, (c_N/\lambda)^{1/(N-1+\alpha)})} ((c_N/\lambda) |x-t|^{1-N})^{\gamma/\alpha} dx \\ &\leq C \lambda^{-\gamma/\alpha} \int_0^{(c_N/\lambda)^{1/(N-1+\alpha)}} r^{N-1-(N-1)\gamma/\alpha} dr \leq C \lambda^{-(N+\gamma)/(N-1+\alpha)}, \end{aligned}$$

under the condition $\gamma < \alpha N/(N-1)$. Then Lemma 2.4 applies if $\gamma > \alpha - 1$, that is $\gamma > 0$ in case $\alpha = 1$. Now assume $\alpha = \gamma = 0$. Then we get directly

$$m_\lambda(t) \leq \int_{B(t, (c_N/\lambda)^{1/(N-1)})} dx \leq C \lambda^{-N/(N-1)}.$$

Hence the functions Φ/ρ and

$$R_i(x) = \int_{\partial\Omega} \frac{\partial \mathcal{G}(x, y)}{\partial x_i} d\varphi(y)$$

lie in $M^{(N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx)$, and satisfy

$$\|\Phi/\rho + |R_1| + \dots + |R_N|\|_{M^{(N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx)} \leq C' \int_{\Omega} \rho^\alpha d|\varphi|.$$

In order to obtain (2.30) and (2.31), it remains to prove that $\partial G(\varphi)/\partial x_i = R_i$ in $\mathcal{D}'(\Omega)$. The result is true when $\varphi \in L^\infty(\Omega)$: in that case, following the proof of [19], Lemma 4.1, we have $G(\varphi) \in C^1(\Omega)$ and $\partial G(\varphi)/\partial x_i = R_i$ in Ω . In the general case where $\varphi \in \mathcal{M}^+(\Omega)$ with $\int_{\Omega} \rho^\alpha d\varphi < +\infty$, we consider a sequence of nonnegative functions $f_n \in L^\infty(\Omega)$, bounded in $L^1(\Omega, \rho^\alpha dx)$, converging weakly to φ . Then the sequence $(G(f_n))$ converges in $L^1(\Omega)$ to $G(\varphi)$ from 2.10. And $(\partial G(f_n)/\partial x_i)$ converges in $L^1(\Omega, \rho dx)$ to R_i . Hence $\partial G(\varphi)/\partial x_i = R_i$ in $\mathcal{D}'(\Omega)$, and in fact in $L^1_{loc}(\Omega)$. ■

Remark 2.3. As a consequence, we get estimates of Ψ and Φ in weighted Sobolev spaces. Recall that for any $k > 1$, and any real γ ,

$$W^{1,k}(\Omega, \rho^\gamma dx) = \{v \in L^k(\Omega, \rho^\gamma dx) \mid |\nabla v| \in L^k(\Omega, \rho^\gamma dx)\}$$

endowed with the norm

$$\|v\|_{W^{1,k}(\Omega, \rho^\gamma dx)} = \|v\|_{L^k(\Omega, \rho^\gamma dx)} + \| |\nabla v| \|_{L^k(\Omega, \rho^\gamma dx)},$$

and $W_0^{1,k}(\Omega, \rho^\gamma dx)$ is the closure of $\mathcal{D}(\Omega)$ in $W^{1,k}(\Omega, \rho^\gamma dx)$. From [16] and [6], it is also given by

$$W_0^{1,k}(\Omega, \rho^\gamma dx) = \{u \in L^k(\Omega, \rho^{\gamma-k} dx) \mid |\nabla u| \in L^k(\Omega, \rho^\gamma dx)\}$$

if $k \neq \gamma + 1$, and

$$W_0^{1,k}(\Omega, \rho^\gamma dx) = \{u \in L^k(\Omega, \rho^{\gamma-k}(\ln(R/\rho))^{-k} dx) \mid |\nabla u| \in L^k(\Omega, \rho^\gamma dx)\}$$

if $k = \gamma + 1$, where $R > \max(e^2, \text{diam } \Omega)$. And

$$W_0^{1,k}(\Omega, \rho^\gamma dx) = W^{1,k}(\Omega, \rho^\gamma dx) \quad \text{if } \gamma + 1 \leq 0 \text{ or } \gamma + 1 > k.$$

Then one verifies that, for any $\mu \in \mathcal{M}(\partial\Omega)$,

$$\Psi = P(\mu) \in W^{1,s}(\Omega, \rho^\gamma dx) \quad (2.47)$$

for any $\gamma > 0$ and $s \in [1, (N + \gamma)/N]$. And for any $\alpha \in [0, 1]$ and any $\varphi \in \mathcal{M}(\Omega)$ such that $\int_\Omega \rho^\alpha d|\varphi| < +\infty$,

$$\Phi = G(\varphi) \in W_0^{1,s}(\Omega, \rho^\gamma dx) \quad (2.48)$$

for any $\gamma \in [0, N\alpha/(N - 1))$ if $\alpha \in (0, 1)$, any $\gamma \in (0, N/(N - 1))$ if $\alpha = 1$, and $\gamma = 0$ if $\alpha = 0$, and for any $s \in [1, (N + \gamma)/(N - 1 + \alpha))$. And P and G map bounded subsets into bounded sets in those spaces. For any measure $\varphi \in \mathcal{M}(\Omega)$, one finds again the well-known result $\Phi = G(\varphi) \in W_0^{1,s}(\Omega)$ for any $s \in [1, N/(N - 1))$.

If $\alpha \in (0, 1)$, we can improve the estimates (2.26) and (2.30) by using *interpolation in weighted spaces*. These results will not be used in the sequel, but they deserve to be mentionned.

Theorem 2.7 *Assume that $\alpha \in (0, 1)$. Then for any $\varphi \in \mathcal{M}(\Omega)$ such that $\int_\Omega \rho^\alpha d|\varphi| < +\infty$, and any $N \geq 3$,*

$$G(\varphi) \in L^{(N+\beta)/(N-2+\alpha)}(\Omega, \rho^\beta dx) \quad (2.49)$$

for any $\beta \in (-N/(N - 1 + \alpha), \alpha N/(N - 2))$, and

$$\|G(\varphi)\|_{L^{(N+\beta)/(N-2+\alpha)}(\Omega, \rho^\beta dx)} \leq C \int_\Omega \rho^\alpha d|\varphi|, \quad (2.50)$$

And for any $N \geq 2$,

$$|\nabla G(\varphi)| + G(\varphi)/\rho \in L^{(N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx) \quad (2.51)$$

for any $\gamma \in [0, \alpha N/(N - 1))$, and

$$\| |\nabla G(\varphi)| + G(\varphi)/\rho \|_{L^{(N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx)} \leq C' \int_\Omega \rho^\alpha d|\varphi|, \quad (2.52)$$

hence

$$G(\varphi) \in W_0^{1, (N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx), \quad (2.53)$$

for any $\gamma \in [0, \alpha N/(N - 1))$, and

$$\|G(\varphi)\|_{W_0^{1, (N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx)} \leq C \int_\Omega \rho^\alpha d|\varphi|. \quad (2.54)$$

Proof. For a given $k > 0$, and any $\alpha_1, \alpha_2 \in [0, 1]$ and $\beta_{i,k} > k + \alpha_i - N$, $\gamma_i > \alpha_i - 1$, for $i = 1, 2$, and any $\theta \in (0, 1)$ we can verify that the spaces of interpolation are given by

$$\begin{aligned} [L^1(\Omega, \rho^{\alpha_1} dx), L^1(\Omega, \rho^{\alpha_2} dx)]_\theta &= L^1(\Omega, \rho^\alpha dx), \\ [M^{(N+\beta_1)/(k+\alpha_1)}(\Omega, \rho^{\beta_1} dx), M^{(N+\beta_2)/(k+\alpha_2)}(\Omega, \rho^{\beta_2} dx)]_\theta &= M^{(N+\beta)/(k+\alpha)}(\Omega, \rho^\beta dx), \end{aligned}$$

where α, β are given by the relations

$$\alpha = (1 - \theta)\alpha_1 + \theta\alpha_2, \quad \frac{1}{p} = \frac{1 - \theta}{p_1} + \frac{\theta}{p_2}, \quad (2.55)$$

$$p_i = (N + \beta_i)/(k + \alpha_i) \quad \text{and} \quad p = (N + \beta)/(k + \alpha), \quad (2.56)$$

From the Marcinkiewicz theorem, if a transformation maps continuously $L^1(\Omega, \rho^{\alpha_i} dx)$ into $M^{(N+\beta_i)/(k+\alpha_i)}(\Omega, \rho^{\beta_i} dx)$ for $i = 1, 2$, it also maps continuously $L^1(\Omega, \rho^\alpha dx)$ into $L^{(N+\beta)/(k+\alpha)}(\Omega, \rho^\beta dx)$, see [29]. Let us show that the estimates (2.49) and (2.51) can be obtained by interpolation of the estimates (2.26), (2.27) and (2.30), (2.31) for $\alpha_1 = 0$ and $\alpha_2 = 1$, *with the exception of the case $\beta = 0$ for (2.51)*. First take $k = N - 2$, and observe that

$$\beta_i \in (-N/(N - 1 + \alpha_i), \alpha_i N/(N - 2)) \Leftrightarrow 1/p_i \in ((N - 2)/N, (N - 1 + \alpha_i)/N), \quad (2.57)$$

so that from (2.55) and (2.56), if $\beta_i \in (-N/(N - 1 + \alpha_i), \alpha_i N/(N - 2))$, then

$$\beta \in (-N/(N - 1 + \alpha), \alpha N/(N - 2)). \quad (2.58)$$

Reciprocally, for any $\alpha \in (0, 1)$ and β satisfying (2.58), taking $\alpha_1 = 0$ and $\alpha_2 = 1$ and defining p by (2.56) with $k = N - 2$, we set

$$p_1 = p_2 = p \quad \text{if } 1/p \leq (N - 2 + \alpha)/N,$$

$$1/p_1 = 1/p - \alpha/N, \quad 1/p_2 = 1/p + (1 - \alpha)/N, \quad \text{if } 1/p > (N - 2 + \alpha)/N.$$

Then p_1 and p_2 satisfy (2.57), and we can interpolate between these values, with β_1, β_2 given by (2.56). Thus G maps continuously $L^1(\Omega, \rho^\alpha dx)$ into $L^{(N+\beta)/(N-2+\alpha)}(\Omega, \rho^\beta dx)$ and (2.50) follows on $L^1(\Omega, \rho^\alpha dx)$. Now take $k = N - 1$, replace β_1, β_2, β by $\gamma_1, \gamma_2, \gamma$ in (2.56): now

$$p_i = (N + \gamma_i)/(N - 1 + \alpha_i) \quad \text{and} \quad p = (N + \gamma)/(N - 1 + \alpha). \quad (2.59)$$

Observe that

$$\gamma_1 = 0 \Leftrightarrow 1/p_1 = (N - 1)/N, \quad (2.60)$$

$$\gamma_2 \in [0, N/(N - 1)) \Leftrightarrow 1/p_2 \in ((N - 1)/N, 1], \quad (2.61)$$

so that from (2.55) and (2.56), if $\gamma_2 \in (0, N/(N-1))$,

$$\gamma \in (0, \alpha N/(N-1)). \quad (2.62)$$

Reciprocally, for any $\alpha \in (0, 1)$ and γ satisfying (2.62), taking $\alpha_1 = 0$ and $\alpha_2 = 1$ and defining p by (2.59), we set

$$1/p_1 = (N-1)/N, \quad 1/p_2 = (1/p - (1-\alpha)/p_1)/\alpha.$$

Then p_1 and p_2 satisfy (2.60), (2.61), and we can interpolate between these values, with γ_1, γ_2 given by (2.59). Hence (2.51) and (2.52) follow on $L^1(\Omega, \rho^\alpha dx)$ when $\gamma \neq 0$. In case $\gamma = 0$, we interpolate between $\alpha_1 = \alpha - \varepsilon$ and $\alpha_2 = \alpha + \varepsilon$ for $\varepsilon > 0$ small enough, with $\gamma_1 = \gamma_2 = 0$, and get again (2.51) and (2.52).

Now consider any $\varphi \in \mathcal{M}(\Omega)$ such that $\int_\Omega \rho^\alpha d|\varphi| < +\infty$. Then there exists a bounded sequence of functions $f_n \in L^1(\Omega, \rho^\alpha dx)$ converging weakly to φ . The sequence $(G(f_n))$ is bounded in $W_0^{1,s}(\Omega)$ for any $s \in [1, N/(N-1+\alpha))$, from (2.48). After an extraction it converges to some function Φ strongly in $L^s(\Omega)$ and a.e. in Ω . Then Φ is a weak solution of problem (2.25), hence $\Phi = G(\varphi)$. Moreover $(G(f_n))$ is bounded in $L^{(N+\beta)/(N-2+\alpha)}(\Omega, \rho^\beta dx)$ for any $\beta \in (-N/(N-1+\alpha), \alpha N/(N-2))$. And $(|\nabla G(f_n)|)$ is bounded in $L^{(N+\gamma)/(N-1+\alpha)}(\Omega, \rho^\gamma dx)$ for any $\gamma \in [0, \alpha N/(N-1))$. Since those spaces are reflexive, we get (2.49) and (2.50), (2.51) and (2.52) by going to the weak limit after a new extraction. Then (2.53) and (2.54) follow. ■

Remark 2.4 Let us mention that the result $\Phi \in W_0^{1,s}(\Omega)$ with $s = N/(N-1+\alpha)$ can be proved by duality, see [12]. Notice that the value of s given in [12] is not correct, due to a small error in the parameters of the Sobolev injection.

Remark 2.5 Assume $N \geq 2$. From (2.48), we deduce that

$$G \text{ is compact from } L^1(\Omega, \rho^\alpha dx) \text{ into } L^p(\Omega, \rho^\beta dx)$$

for any $\alpha \in (0, 1]$, $\beta \in (-N/(N+\alpha-1), N\alpha/(N-2))$ or $\alpha = \beta = 0$, and $p \in [1, (N+\beta)/(N+\alpha-2))$ and $p > -\beta$. It comes from the compactness of the Sobolev injection

$$W_0^{1,s}(\Omega, \rho^\gamma dx) \subset L^p(\Omega, \rho^\beta dx)$$

when $1 \leq s \leq p < +\infty$ and $N/p - N/s + 1 > 0$ and $(N+\beta)/p - (N+\gamma)/s + 1 > 0$, with $\gamma + 1 \neq s$, see [27]. In the case $\alpha = 1$ and $\beta = 0$, we find again a result cited in [10].

2.3 Application to the problems 1.1 and 1.4

Combining Theorems 2.5 and 2.6, we deduce regularity results for the problem (1.4). In particular, taking $\alpha = 1$, we get the following:

Corollary 2.8 *For any $\varphi \in \mathcal{M}(\Omega)$ such that $\int_{\Omega} \rho d|\varphi| < +\infty$ and any $\mu \in \mathcal{M}(\partial\Omega)$, the solution U of problem (1.4) satisfies*

$$\begin{cases} U \in M^{(N+\beta)/(N-1)}(\Omega, \rho^{\beta} dx) & \text{for any } \beta \in (-1, N/(N-2)), \text{ if } N \geq 3, \\ U \in L^k(\Omega, \rho^{\beta} dx) & \text{for any } \beta \in (-1, +\infty) \text{ and } k \in [1, 2+\beta), \text{ if } N = 2, \end{cases} \quad (2.63)$$

$$|\nabla U| \in M^{(N+\gamma)/N}(\Omega, \rho^{\gamma} dx), \quad \text{for any } \gamma \in (0, N/(N-1)), \text{ if } N \geq 2, \quad (2.64)$$

(hence $U \in W_0^{1,s}(\Omega, \rho^{\gamma} dx)$ for any $\gamma \in (0, N/(N-1))$ and $s \in [1, (N+\gamma)/N)$). And in any case,

$$\|U\|_{M^{(N+\beta)/(N-1)}(\Omega, \rho^{\beta} dx)} \leq C \left[\int_{\Omega} \rho d|\varphi| + |\mu|(\partial\Omega) \right], \text{ if } N \geq 3, \quad (2.65)$$

$$\| |\nabla U| \|_{M^{(N+\gamma)/N}(\Omega, \rho^{\gamma} dx)} \leq C \left[\int_{\Omega} \rho d|\varphi| + |\mu|(\partial\Omega) \right], \text{ if } N \geq 2. \quad (2.66)$$

This gives an interior regularity result for problem (1.5):

Corollary 2.9 *If $1 < q < (N+1)/(N-1)$, then any solution u of (1.1) is a classical solution in Ω .*

Proof. Applying Corollary 2.8 to u we get in particular $u \in M^{(N+1)/(N-1)}(\Omega, \rho dx)$ if $N \geq 3$ (and $u \in M^{3-\varepsilon}(\Omega, \rho dx)$ if $N = 2$). Then $u^q \in L_{loc}^{k_0}(\Omega)$ for some $k_0 > 1$, since $q < (N+1)/(N-1)$. If $N = 2$, then from Schauder estimates, $u \in C^{\infty}(\Omega)$. In case $N \geq 3$ and $k_0 < N/2$, we can make a usual bootstrapp: from the L^p regularity theory, $u \in W_{loc}^{2,k_0}(\Omega)$, hence from the Sobolev injection $u^q \in L_{loc}^{k_1}(\Omega)$ for $k_1 = N k_0 / q (N - 2k_0) > k_0$, since $q < N/(N-2)$. By induction $u^q \in L_{loc}^{k_n}(\Omega)$ for

$$k_n = N k_{n-1} / q (N - 2 k_{n-1}) > k_{n-1}, \quad (2.67)$$

till $k_n < N/2$. But if $k_n < N/2$ for any $n \in \mathbb{N}$, then $k_n \rightarrow \ell = N(q-1)/2q < 1$, which is impossible. Then changing slightly k_0 if necessary, we find some $n_0 \in \mathbb{N}$ such that $k_{n_0} > N/2$, hence $u \in C^{\infty}(\Omega)$. ■

3 Estimate of $G(P^q(\mu))$

Now we assume that $1 < q < (N+1)/(N-1)$, and we prove Theorem 1.1. First observe that for any $\mu \in \mathcal{M}^+(\partial\Omega)$, we have $P(\mu) \in M^{(N+1)/(N-1)}(\Omega, \rho dx)$ from Theorem 2.5. In particular,

$$P^q(\mu) \in L^1(\Omega, \rho dx)$$

since $q < (N + 1)/(N - 1)$, hence $G(P^q(\mu))$ is well defined and lies in $L^1(\Omega)$ from Corollary 2.2. And $P(\mu) \in C^0(\Omega)$, since \mathcal{P} is continuous, hence also $G(P^q(\mu)) \in C^0(\Omega)$.

We first consider the case where $\mu = \delta_a$ is a Dirac mass at a point a of $\partial\Omega$, hence $P(\mu)(.) = P(\delta_a)(.) = \mathcal{P}(., a)$. Here we can give a more precise estimate near the point a :

Theorem 3.1 *Assume that $1 < q < (N + 1)/(N - 1)$. Let $a \in \partial\Omega$, and let $W = G(P^q(\delta_a))$ be the solution of*

$$\begin{cases} -\Delta W = \mathcal{P}^q(., a) & \text{in } \Omega, \\ W = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.1)$$

Then there exists a constant $C = C(N, \Omega, q)$ such that

$$W(x) \leq C \mathcal{P}(x, a) |x - a|^{N+1-(N-1)q} \quad \text{in } \Omega. \quad (3.2)$$

Proof. From (2.8), we can majorize $\mathcal{P}^q(., a)$ by

$$\mathcal{P}^q(x, a) \leq c_N^q \rho^q(x) |x - a|^{-Nq} \leq c_N^q \rho(x) |x - a|^{-1-(N-1)q}.$$

Then for any $x \in \Omega$, from (1.3),

$$W(x) = \int_{\Omega} \mathcal{G}(x, y) \mathcal{P}^q(y, a) dy \leq c_N^q \int_{\Omega} \mathcal{G}(x, y) \rho(y) |y - a|^{-1-(N-1)q} dy.$$

Now from (2.2) and (2.3),

$$W(x) \leq c_N^{q+1} \rho(x) \int_{\Omega} f(x, y) dy,$$

where

$$f(x, y) = |y - a|^{-(N-1)q} |x - y|^{-N} \min(|x - y|, |y - a|),$$

since $\rho(y) \leq |y - a|$. Now we divide Ω in three parts:

$$\Omega_1 = \Omega \cap B(x, |x - a|/2), \quad \Omega_2 = \Omega \cap B(a, |x - a|/2), \quad \Omega_3 = \Omega \setminus (\Omega_1 \cup \Omega_2),$$

and integrate separately on each part. In the sequel C denotes constants which only depend on N, q and Ω . In Ω_1 we have $|x - a| \leq 2|y - a|$, and

$$\begin{aligned} \int_{\Omega_1} f(x, y) dy &\leq \int_{\Omega_1} |y - a|^{-(N-1)q} |x - y|^{1-N} dy \\ &\leq 2^{(N-1)q} |x - a|^{-(N-1)q} \int_{B(x, |x-a|/2)} |x - y|^{1-N} dy \\ &\leq C |x - a|^{1-(N-1)q}. \end{aligned}$$

In Ω_2 we have $|x - a| \leq 2|x - y|$, and

$$\begin{aligned}
\int_{\Omega_2} f(x, y) dy &\leq \int_{\Omega_2} |y - a|^{1-(N-1)q} |x - y|^{-N} dy \\
&\leq 2^N |x - a|^{-N} \int_{B(a, |x-a|/2)} |y - a|^{1-(N-1)q} dy \\
&\leq C |x - a|^{-N} \int_0^{|x-a|/2} r^{N-(N-1)q} dr \leq C |x - a|^{1-(N-1)q},
\end{aligned}$$

since $q < (N + 1)/(N - 1)$.

In Ω_3 , we have $|x - a| \leq 2 \min(|y - a|, |y - x|)$, hence $|y - a| \leq 3|y - x|$. Then we get

$$\begin{aligned}
\int_{\Omega_3} f(x, y) dy &\leq \int_{\Omega_3} |y - a|^{1-(N-1)q} |x - y|^{-N} dy \\
&\leq 3^{-N} \int_{\Omega_3} |y - a|^{1-N-(N-1)q} dy \\
&\leq C \int_{|x-a|/2}^{+\infty} r^{-(N-1)q} dr \\
&\leq C |x - a|^{1-(N-1)q}.
\end{aligned}$$

Then

$$W(x) \leq C \rho(x) |x - a|^{1-(N-1)q},$$

and from the lower estimate of the Poisson kernel (2.8) we deduce (3.2). ■

Now we get to the general case.

Proof of Theorem 1.1 Let $\mu \in \mathcal{M}^+(\partial\Omega)$. We can reduce to the case $\mu(\partial\Omega) = 1$ by linearity of P and G . From (1.2), we have

$$P(\mu)(x) = \int_{\partial\Omega} \mathcal{P}(x, z) d\mu(z) = \int_{\partial\Omega} P(\delta_z)(x) d\mu(z) \quad \text{in } \Omega.$$

Then from the Jensen inequality,

$$P^q(\mu)(x) \leq \int_{\partial\Omega} P^q(\delta_z)(x) d\mu(z) \quad \text{in } \Omega.$$

And from the maximum principle,

$$G(P^q(\mu))(x) \leq G\left(\int_{\partial\Omega} P^q(\delta_z)(x) d\mu(z)\right) = \int_{\partial\Omega} G(P^q(\delta_z))(x) d\mu(z).$$

Hence from (3.2),

$$\begin{aligned} G(P^q(\mu))(x) &\leq C \int_{\partial\Omega} \mathcal{P}(x, z) |x - z|^{N+1-(N-1)q} d\mu(z) \\ &\leq C P(\mu)(x) \quad \text{in } \Omega, \end{aligned}$$

since $N + 1 - (N - 1)q > 0$ and Ω is bounded. ■

4 A priori estimates

Here we study the behaviour of the solutions of (1.5) for a given measure $\mu \in \mathcal{M}^+(\partial\Omega)$. First notice as in [13] that for any $q > 1$, for any solution u of (1.5), $\|u^q\|_{L^1(\Omega, \rho dx)}$ is majorized independently of u : we have the estimate

$$\|u^q\|_{L^1(\Omega, \rho dx)} \leq C (1 + \mu(\partial\Omega)), \quad (4.1)$$

with $C = C(N, q, \Omega)$. Indeed consider a positive eigenvector Φ_1 for the first eigenvalue λ_1 of $(-\Delta)$ with Dirichlet conditions on $\partial\Omega$. Since u is a weak solution of (1.5), we have

$$\int_{\Omega} u (-\Delta\Phi_1) dx = \lambda_1 \int_{\Omega} u \Phi_1 dx = \int_{\Omega} u^q \Phi_1 dx + \int_{\partial\Omega} \frac{\partial\Phi_1}{\partial n} d\mu,$$

hence from Young inequality

$$\int_{\Omega} u^q \Phi_1 dx \leq \frac{1}{2} \int_{\Omega} u^q \Phi_1 dx + (2\lambda_1^q)^{1/(q-1)} \int_{\Omega} \Phi_1 dx + \int_{\partial\Omega} \left| \frac{\partial\Phi_1}{\partial n} \right| d\mu,$$

which implies (4.1), since $C^{-1}\rho \leq \Phi_1 \leq C^{-1}\rho$ in Ω , for some $C = C(N, \Omega) > 0$.

Now we prove Theorem 1.1. We follow the technique of the interior problem, given in [23]. Once we have obtained the estimate (1.12), the proof goes quickly in case $q < N/(N-1)$. The main difficulty comes when $q \geq N/(N-1)$: in that case we really need the precise estimates of $G(\varphi)$ and $P(\mu)$ in Marcinkiewicz weighted spaces, proved in Section 2.2. We begin by the easiest case.

Proof of Theorem 1.1 .

i) The simple case : $q < N/(N-1)$.

Let $\mu \in \mathcal{M}^+(\partial\Omega)$, and let u be any nonnegative solution of (1.5). Let us set

$$u = P(\mu) + v_1 \quad (4.2)$$

where $v_1 = G(u^q)$. Now

$$u \in M^{N/(N-1)}(\Omega) \text{ if } N \geq 3, \quad u \in M^{2-\varepsilon}(\Omega) \text{ if } N = 2, \quad (4.3)$$

from Corollary (2.8). Since $q < N/(N-1)$, $u^q \in L^{k_0}(\Omega)$ for some $k_0 > 1$. Now $u^q \leq 2^{q-1}(P^q(\mu) + v_1^q)$, hence from the maximum principle and from the estimate (1.12),

$$v_1 \leq 2^{q-1}(G(P^q(\mu)) + v_2) \leq C_1(P(\mu) + v_2)$$

where $v_2 = G(v_1^q)$, hence

$$u \leq C'_1(P(\mu) + v_2).$$

By induction for any $n \geq 2$, we can define the solution $v_n = G(v_{n-1}^q)$ of problem

$$\begin{cases} -\Delta v_n = v_{n-1}^q & \text{in } \Omega \\ v_n = 0 & \text{on } \partial\Omega, \end{cases}$$

such that

$$v_n \leq C_n(P(\mu) + v_{n+1}), \quad u \leq C'_n(P(\mu) + v_{n+1}),$$

where C_n, C'_n only depend on N, q, Ω and $\mu(\partial\Omega)$. And $v_n \in L^{k_n}(\Omega)$ with k_n given by (2.67), hence there exists some $n_0 = n_0(N, q)$ such that $v_{n_0} \in C^0(\overline{\Omega})$. Then

$$u \leq C'_{n_0}(P(\mu) + v_{n_0+1}) \quad \text{in } \Omega. \quad (4.4)$$

with $v_{n_0+1} = 0$ on $\partial\Omega$, hence there exists a constant $C_0 > 0$ such that

$$v_{n_0+1}(x) \leq C_0 \rho(x) \quad \text{in } \Omega, \quad (4.5)$$

and C_0 depends on $N, q, \Omega, \mu(\partial\Omega)$ and $\|u^q\|_{L^1(\Omega, \rho dx)}$, from the continuity properties given in Corollary 2.8, hence $C_0 = C_0(N, q, \Omega, \mu(\partial\Omega))$ from (4.1). Then (1.13) follows from (4.4) and (4.5). If $\mu = 0$, then $u \in C^\infty(\overline{\Omega})$ from Schauder estimates.

In case $\mu = \sigma \delta_a$ for some $a \in \partial\Omega$ and $\sigma > 0$, we get more precisely from Theorem 3.1

$$\begin{aligned} v_1 &\leq 2^{q-1}(\sigma^q G(P^q(\delta_a)) + v_2) \\ &\leq C_1 \left[P(\delta_a) |x - a|^{N+1-(N-1)q} + v_2 \right]. \end{aligned}$$

By induction we find

$$\begin{aligned} v_n &\leq C_n (G(P^q(\delta_a)) + v_{n+1}) \\ &\leq C'_n \left[P(\delta_a) |x - a|^{N+1-(N-1)q} + v_{n+1} \right] \end{aligned}$$

and

$$u \leq P(\delta_a) + C'_n \left[P(\delta_a) |x - a|^{N+1-(N-1)q} + v_{n+1} \right].$$

Then we deduce (1.14).

ii) The case: $N/(N-1) \leq q < (N+1)/(N-1)$.

Let $p \geq 2$ be some fixed integer such that

$$1/p < N + 1 - (N - 1)q.$$

Now for any $n \in [0, p]$, let $\beta_n = 1 - n/p \in [0, 1]$. Now we start from the fact that

$$u \in M^{(N+1)/(N-1)}(\Omega, \rho \, dx) \text{ if } N \geq 3, \quad u \in M^{3-\varepsilon}(\Omega, \rho \, dx) \text{ if } N = 2, \quad (4.6)$$

from Corollary 2.8 Let $v_0 = u$, hence

$$v_0^q \in L^{r_0}(\Omega, \rho^{\beta_0} \, dx), \quad \text{with } 1 < r_0 < (N + \beta_0)/(N - 1)q.$$

Here again we define v_1 by (4.2), and $v_1 \leq u$. So that we can define $v_2 = G(v_1^q)$ in $L^1(\Omega)$. From Theorem 2.6, we have, for any $N \geq 2$ and $\varepsilon > 0$ small enough,

$$v_1 \in L^{(N+\beta)/(N-2+\beta_0)-\varepsilon}(\Omega, \rho^\beta \, dx)$$

for any $\beta \in (-1, N/(N - 2))$. Taking $\beta = \beta_1 = 1 - 1/p \in (0, 1)$, we get

$$v_1^q \in L^{r_1}(\Omega, \rho^{\beta_1} \, dx), \quad \text{with } 1 < r_1 < (N + \beta_1)/(N - 2 + \beta_0)q, \quad (4.7)$$

since $N + \beta_1 - (N - 2 + \beta_0)q = N + 1 - (N - 1)q - 1/p > 0$. For any $n \leq p$, assume by induction that $v_{n-1} = G(v_{n-2}^q)$ in $L^1(\Omega)$, and that

$$v_{n-1}^q \in L^{r_{n-1}}(\Omega, \rho^{\beta_{n-1}} \, dx), \quad \text{with } 1 < r_{n-1} < (N + \beta_{n-1})/(N - 2 + \beta_{n-2})q,$$

then we can define $v_n = G(v_{n-1}^q)$ in $L^1(\Omega)$, and we get

$$v_n \in L^{(N+\beta)/(N-2+\beta_{n-1})-\varepsilon}(\Omega, \rho^\beta \, dx)$$

for any $\beta \in (-1, \beta_{n-1}N/(N - 2))$. Taking $\beta = \beta_n \geq 0$, we have $(N + \beta_n) - (N - 2 + \beta_{n-1})q > (n - 1)(q - 1)/p > 0$, hence

$$v_n^q \in L^{r_n}(\Omega, \rho^{\beta_n} \, dx), \quad \text{with } 1 < r_n < (N + \beta_n)/(N - 2 + \beta_{n-1})q. \quad (4.8)$$

Now in case $n = p$, we have $\beta_p = 0$. This proves that $v_p^q \in L^{r_p}(\Omega)$, with $r_p > 1$ and we are reduced to the first case: there exists an integer $n_0 = n_0(N, q)$ such that $v_{n_0+p} \in C^0(\overline{\Omega})$. We deduce (1.13) and (1.14) as above. ■

In this proof we have used the estimate (1.12). In fact it is not really needed for getting a priori estimates, since we require that the problem admits a solution: the existence assumption in turn implies a condition of type (??). Adapting the arguments of [8] for the interior nonhomogeneous problem

$$\begin{cases} -\Delta u = u^q + f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

with $f > 0$, we get the following:

Lemma 4.1 *Let $q > 1$ and $\mu \in \mathcal{M}^+(\partial\Omega)$. If the problem (1.5) admits a solution, then*

$$G(P^q(\mu)) \leq \frac{1}{q-1}P(\mu) \quad \text{in } \Omega. \quad (4.9)$$

Proof. We can assume $\mu \neq 0$. For any $v, w \in C^2(\Omega)$ with v positive and harmonic, and any concave function F of class C^2 on the closure of the range of w/v , we have

$$-\Delta [v F(w/v)] \geq F'(w/v) (-\Delta w) \quad \text{in } \Omega. \quad (4.10)$$

Suppose that problem (1.5) admits a solution u . Then we apply (4.10) with $v = P(\mu)$ and $w = u \geq v$, and

$$F(s) = (1 - s^{1-q}) / (q-1) \quad \text{on } [1, +\infty).$$

It comes

$$P^q(\mu) = [u/P(\mu)]^{-q} u^q \leq -\Delta [P(\mu) F[u/P(\mu)]]$$

and

$$G(P^q(\mu)) \leq P(\mu) F[u/P(\mu)] \leq \frac{1}{q-1}P(\mu),$$

from the maximum principle. ■

Remark 4.1 In the supercritical case $q \geq (N+1)/(N-1)$, any solution u of (1.1), such that $u \in C(\bar{\Omega} \setminus \{a\})$ and $u = 0$ on $\partial\Omega \setminus \{a\}$, satisfies necessarily $u = 0$ on $\partial\Omega$, from Corollary 2.3. Considering the known behaviours for the problems (1.7) and (1.6), one can ask if an estimate of the type

$$u(x) \leq C \rho(x) |x - a|^{-(q+1)/(q-1)}$$

is true near the point a , at least if $q < (N+1)/(N-3)$. The question is entirely open.

5 Existence results

Here we study the existence of solutions of problem (1.15). It is based on the estimate of $G(P^q(\mu))$, which gives supersolutions:

Proof of Theorem 1.2

First step: existence of solutions for small σ . Let $\mu \in \mathcal{M}^+(\partial\Omega)$ with $\mu(\partial\Omega) = 1$ and $\sigma > 0$. The function $\sigma P(\mu)$ is a subsolution of (1.15). We search a supersolution of (1.15) of the form

$$y = \sigma P(\mu) + a G[P^q(\sigma \mu)]$$

with $a > 0$, and

$$a P^q(\sigma \mu) \geq y^q.$$

Since $q < (N+1)/(N-1)$, from Theorem 1.1, there exists a constant $K = K(N, \Omega, q)$ such that

$$y \leq \sigma(1 + a\sigma^{q-1}K) P(\mu) \quad \text{in } \Omega, \quad (5.1)$$

and y is a supersolution as soon as $a^{1/q} \geq 1 + a\sigma^{q-1}K$. As a consequence, taking the best value $a = (q/(q-1))^q$, if

$$\sigma \leq (qK)^{-1/(q-1)}(q-1)/q, \quad (5.2)$$

then \mathcal{S}_σ has a solution.

Second step: interval of existence. Let

$$\Lambda = \{\sigma > 0 \mid \mathcal{S}_\sigma \text{ has a solution}\} \quad \text{and} \quad \sigma^* = \sup \Lambda.$$

Then from (4.9),

$$(\sigma^*)^{q-1} G(P^q(\mu)) \leq \frac{1}{q-1} P(\mu) \quad \text{in } \Omega, \quad (5.3)$$

hence σ^* is finite. For any $\sigma \in \Lambda$, \mathcal{S}_σ has a solution u_σ . For any $\tau \in [0, \sigma)$, u_σ is a supersolution of (1.1) such that $u_\sigma \geq \tau P(\mu)$, hence \mathcal{S}_τ has a solution $u_\tau \leq u_\sigma$. Then Λ is an interval. At last, let us show that \mathcal{S}_{σ^*} has a solution: let (σ_n) be an increasing sequence with limit σ^* . Now u_{σ_n} is a weak solution of \mathcal{S}_{σ_n} ; we use as a test function the unique solution $\xi > 0$ of problem $\xi = G(\xi^{1/q})$, introduced in [8], and get

$$\int_{\Omega} u_{\sigma_n} (-\Delta \xi) dx = \int_{\Omega} u_{\sigma_n} \xi^{1/q} dx = \int_{\Omega} u_{\sigma_n}^q \xi dx - \sigma_n \int_{\partial\Omega} \frac{\partial \xi}{\partial n} d\mu. \quad (5.4)$$

And $\partial \xi / \partial n < 0$, hence

$$\int_{\Omega} u_{\sigma_n}^q \xi dx \leq \frac{1}{q} \int_{\Omega} u_{\sigma_n}^q \xi dx + \frac{q-1}{q} |\Omega|,$$

so that $(u_{\sigma_n}^q)$ is bounded in $L^1(\Omega, \rho dx)$. And $u_{\sigma_n} \leq \sigma^* P(\mu)$ on $\partial\Omega$. Now (u_{σ_n}) is bounded in $M^{N/(N-1)}(\Omega) \cap M^{(N+1)/(N-1)}(\Omega, \rho dx)$, from Corollary 2.8. Then we can go to the limit in the weak formulation of \mathcal{S}_{σ_n} , and construct a weak solution of \mathcal{S}_{σ^*} . \blacksquare

Remark 5.1 Taking $K = \|G(P^q(\mu))/P(\mu)\|_{L^\infty(\Omega)}$, we can estimate σ^* from (5.2) and (5.3):

$$\sigma^* \in [(qK)^{-1/(q-1)}(q-1)/q, ((q-1)K)^{-1/(q-1)}].$$

Remark 5.2 Now assume $q \geq (N+1)/(N-1)$. We shall say that a measure $\mu \in \mathcal{M}^+(\partial\Omega)$ with $\mu(\partial\Omega) = 1$ is *admissible* if

$$P^q(\mu) \in L^1(\Omega) \text{ and } \mu \text{ satisfies the condition (??) for some } K > 0.$$

Then in the same way Theorem 1.2 applies to any admissible measure. Moreover, for such an admissible measure, following the techniques of [8], for any $\sigma \in [0, \sigma^*)$, we can construct a solution u_σ of \mathcal{S}_σ satisfying the a priori estimate

$$\sigma P(\mu) \leq u_\sigma \leq C P(\mu) \quad \text{in } \Omega, \quad (5.5)$$

for some constant $C = C(\sigma)$. Indeed let $\tau \in (\sigma, \sigma^*)$ such that \mathcal{S}_τ admits a solution u_τ . Let us apply the inequality (4.10) with $v = \tau P(\mu)$ and $w = u_\tau \geq v$, and

$$F(s) = s \left(1 + \varepsilon s^{q-1}\right)^{-1/(q-1)} \quad \text{and} \quad \varepsilon = (\tau/\sigma)^{q-1} - 1, \quad \text{on } [1, +\infty),$$

so that $F(1) = \sigma/\tau$ and $F'(s) = F^q(s)/s^q$:

$$-\Delta[v F(u_\tau/v)] \geq F'(u_\tau/v) (-\Delta u_\tau) = [v F(u_\tau/v)]^q \quad \text{in } \Omega.$$

Hence $z = v F(u_\tau/v)$ is a supersolution of (1.1). Then

$$z \geq \sigma P(\mu) \quad \text{in } \Omega, \quad \text{and } z = \sigma \mu \quad \text{on } \partial\Omega$$

and \mathcal{S}_σ has a solution $u_\sigma \leq z \leq \varepsilon^{-1/(q-1)} \tau P(\mu)$, so that u_σ satisfies (5.5). Now for any $\tau \in (\sigma, \sigma^*)$, \mathcal{S}_τ admits a solution u_τ . Choosing $\tau = (\sigma + \sigma^*)/2$, we deduce that u_σ satisfies (5.5) with $C(\sigma) = \sigma^* [(\sigma + \sigma^*)/2\sigma]^{q-1} - 1]^{-1/(q-1)}$. At last considering as above an increasing sequence (σ_n) with limit σ^* , we prove that \mathcal{S}_{σ^*} admits a solution u_{σ^*} .

An open question is to describe precisely those admissible measures.

References

- [1] H. Amann and P. Quittner, *Elliptic boundary value problems involving measures: existence, regularity and multiplicity*, preprint.
- [2] P. Benilan, H. Brézis and M.G. Crandall, *A semilinear elliptic equation in $L^1(\mathbb{R}^N)$* , Ann. Scuol. Norm. Sup. Pisa, 2 (1975), 523-555.
- [3] M.F. Bidaut-Véron, *Local and global behaviour of solutions of quasilinear equations of Emden-Fowler type*, Arc. Rat. Mech. Anal., 107 (1989), 293-324.
- [4] M.F. Bidaut-Véron, *Local behaviour of solutions of a class of nonlinear elliptic systems*, Advances in Diff. Equ., to appear.
- [5] M.F. Bidaut-Véron and L. Véron, *Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations*, Invent. Math. 106 (1991), 489-539.

- [6] P. Bolley and J. Camus, *Quelques résultats sur les espaces de Sobolev à poids*, Publ. Séminaires Math. Univ. Rennes (1968-69), 1-70.
- [7] H. Brézis, *Une équation semilinéaire avec conditions aux limites dans L^1* , unpublished work.
- [8] H. Brézis and X. Cabré, *Some simple nonlinear PDE's without solutions*, Boll. Unione Mat. Italiana, 8 (1998), 223-262.
- [9] H. Brézis and P.L. Lions, *A note on isolated singularities for linear elliptic equations*, Math. Anal. Appl. Adv. Math., Suppl. Stud. 7A (1981), 263-266.
- [10] X. Cabré and Y. Martel, *Weak eigenfunctions for the linearization of extremal elliptic problems*, J. Funct. Anal., to appear.
- [11] L.A. Caffarelli, B. Gidas and J. Spruck, *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*, Comm. Pure Appl. Math., 42 (1989), 271-297.
- [12] R. Dautray and J.L. Lions, *Analyse mathématique et calcul numérique*, Masson ed. (1987).
- [13] D.G. De Figuereido, P.L. Lions and R.D. Nussbaum, *A priori estimates and existence of positive solutions of semilinear elliptic equations*, J. Math. Pures et Appl., 61 (1982), 41-63.
- [14] E.B. Dynkin, *A probabilistic approach to one class of nonlinear differential equations*, Prob. Th. Rel. Fields, 89 (1991), 89-115.
- [15] E.B. Dynkin and S.E. Kuznetsov, *Superdiffusion and removable singularities for quasilinear partial differential equations*, Comm. Pure Appl. Math. 49 (1996), 125-176.
- [16] G. Geymonat and P. Grisvard, *Problemi ai limiti lineari ellittici negli spazi di Sobolev con peso*, Le Matematiche, 22, 2 (1967), 1-38.
- [17] B. Gidas and J. Spruck, *Global and local behavior of positive solutions of nonlinear elliptic equations*, Comm. Pure Appl. Math. 34 (1981), 525-598.
- [18] B. Gidas and J. Spruck, *A priori bounds for positive solutions of nonlinear elliptic equations*, Comm. Part. Diff. Equ., 6 (1981), 883-901.
- [19] D. Gilbarg and N.S. Trudinger, *Elliptic partial differential equations of second order*, Springer-Verlag, Berlin Heidelberg New-York Tokyo (1983).

- [20] A. Gmira and L. Véron, *Boundary singularities of solutions of some nonlinear elliptic equations*, Duke Math. Journal, 64 (1991), 271-324.
- [21] S.G. Krantz, *Function theory of several complex variables*, 2nd ed., Wadsworth, Belmont (1992).
- [22] J.F. Le Gall, *Solutions positives de $\Delta u = u^2$ dans le disque unité*, C.R. Acad. Sci. 317 série I (1993), 873-878.
- [23] P.L. Lions, *Isolated singularities in semilinear problems*, J. Diff. Equ., 38 (1980), 441-450.
- [24] M. Marcus and L.Véron, *The boundary trace of positive solutions of semilinear elliptic equations: I The subcritical case*, Arch. Rat. Mech. and Anal., to appear.
- [25] M. Marcus and L.Véron, *The boundary trace of positive solutions of semilinear elliptic equations: The supercritical case*, J. Math. Pures Appl., 77 (1998), 481-524.
- [26] V.G. Maz'ja, *Beurling's theorem on a minimum principle for positive harmonic functions*, J. Sov. Math. 4 (1975), 367-379.
- [27] B. Opic and A. Kufner, *Hardy type inequalities*, Pitman Research in Math. Series, Longman Scientific Technical (1990).
- [28] J. Serrin, *Isolated singularities of solutions of quasilinear equations*, Acta Math. 1134 (1964), 247-302.
- [29] H. Triebel, *Interpolation theory, Function Spaces, Differential Equations*, North-Holland (1978).
- [30] L. Véron, *Singularities of solutions of second order quasilinear equations*, Pitman Research Notes in Math., Longman Sci. & Tech.353 (1996).
- [31] L. Véron, *Singular solutions of some nonlinear elliptic equations*, Nonlinear Anal. 5 (1981), 225-242.
- [32] L. Vivier, *Thèse de Doctorat*, Tours (1998).
- [33] K.O. Widman, *Inequalities for the Green function and boundary continuity of the gradient of solutions of elliptic differential equations*, Math. Scand. 21 (1967), 17-37.