

Brownian motion in cones: algebraic and analytic approaches

$\chi_\lambda = \frac{\sum_{w \in W} \varepsilon(w) e^{w \cdot (\lambda + \rho)}}{\sum_{w \in W} \varepsilon(w) e^{w \cdot \rho}}$ Toulouse, November 6-8, 2017

Speakers:

$$\mathcal{P}B_t = B_t - 2 \inf_{0 \leq s \leq t} B_s$$

Julien Barré (Univ. Orléans & IUF)

Philippe Bougerol (Univ. Pierre et Marie Curie)

Guillaume Cébron (Univ. Toulouse)

Emmanuel Chasseigne (Univ. Tours)

Nizar Demni (Univ. Rennes)

Sandro Franceschi (Univ. Pierre et Marie Curie)

Cédric Lecouvey (Univ. Tours)

Antoine Lejay (Inria Nancy)

Pierre-Loïc Méliot (Univ. Paris-Sud)

Laurent Miclo (CNRS & Univ. Toulouse)

Gregory Schehr (CNRS & Univ. Paris-Sud)

Organizers:

François Chapon, Reda Chhaibi, Vincent Perrollaz, Kilian Raschel

Funding:

CNRS PEPS project "Analytic and combinatorial approaches of Brownian motion in a cone".

CNRS PEPS project "Quantum random walks and quantum groups", the IUF, the federation Denis Poisson and the Institut de Mathématiques de Toulouse

