

## EXTINCTION PROBABILITIES FOR A DISTYLOUS PLANT POPULATION MODELED BY AN INHOMOGENEOUS RANDOM WALK ON THE POSITIVE QUADRANT\*

PAULINE LAFITTE-GODILLON<sup>†</sup>, KILIAN RASCHEL<sup>‡</sup>, AND VIET CHI TRAN<sup>§</sup>

**Abstract.** In this paper, we study a flower population in which self-reproduction is not permitted. Individuals are diploid, that is, each cell contains two sets of chromosomes, and distylous, that is, two alleles,  $A$  and  $a$ , can be found at the considered locus  $S$ . Pollen and ovules of flowers with the same genotype at locus  $S$  cannot mate. This prevents the pollen of a given flower from fecundating its own stigmata. Only genotypes  $AA$  and  $Aa$  can be maintained in the population, so that the latter can be described by a random walk in the positive quadrant whose components are the number of individuals of each genotype. This random walk is not homogeneous, and its transitions depend on the location of the process. We are interested in the computation of the extinction probabilities, as extinction happens when one of the axes is reached by the process. These extinction probabilities, which depend on the initial condition, satisfy a doubly indexed recurrence equation that cannot be solved directly. Our contribution is twofold: on the one hand, we obtain an explicit, though intricate, solution through the study of the PDE solved by the associated generating function. On the other hand, we provide numerical results comparing stochastic and deterministic approximations of the extinction probabilities.

**Key words.** inhomogeneous random walk on the positive quadrant, boundary absorption, transport equation, method of characteristics, self-incompatibility in flower populations, extinction in diploid population with sexual reproduction

**AMS subject classifications.** 60G50, 60J80, 35Q92, 92D25

**DOI.** 10.1137/120864258

**1. Introduction.** We consider the model of a flower population without pollen limitation introduced by Billiard and Tran [4]. The flower reproduction is sexual: plants produce pollen that may fecundate the stigmata of other plants. We are interested in self-incompatible reproduction, where an individual can reproduce only with compatible partners. In particular, self-incompatible reproduction prevents the fecundation of a plant's stigmata by its own pollen. Each plant is diploid and characterized by the two alleles that it carries at the locus  $S$ , which determine the possible types of partners with whom the plant may reproduce (as it encodes the recognition proteins present on the pollen and stigmata of the plant). We consider the distyle case with only two possible alleles,  $A$  or  $a$ . The plants thus can have genotypes  $AA$ ,  $Aa$ , or  $aa$ . The only interesting case is when  $A$  is dominant over  $a$  (see [4]), and we restrict our analysis to that case in this work. Then, the phenotype, i.e., the type of proteins carried by the pollen and stigmata, of individuals with genotypes  $AA$  (resp.,  $Aa$  and

---

\*Received by the editors January 31, 2012; accepted for publication (in revised form) December 5, 2012; published electronically March 19, 2013.

<http://www.siam.org/journals/siap/73-2/86425.html>

<sup>†</sup>Ecole Centrale de Paris, 92295 Châtenay-Malabry Cédex, France, and EPI SIMPAF, INRIA Lille Nord-Europe, 59650 Villeneuve d'Ascq, France (pauline.lafitte@ecp.fr). This author's work was supported by a "projet BQR Emergent" of USTL.

<sup>‡</sup>CNRS and Laboratoire de Mathématiques et Physique Théorique, Université de Tours, 37200 Tours, France (kilian.raschel@lmpt.univ-tours.fr).

<sup>§</sup>Laboratoire Paul Painlevé, Université des Sciences et Technologies Lille 1, 59655 Villeneuve d'Ascq Cédex, France (chi.tran@math.univ-lille1.fr). This author's work was supported by ANR MANEGE (ANR-09-BLAN-0215), by the "Chaire Modélisation Mathématique et Biodiversité of Veolia Environnement - Ecole Polytechnique - Museum National d'Histoire Naturelle - Fondation X," and by a "projet BQR Emergent" of USTL.




FIG. 1.1. (a) Transition rates for the continuous-time pure-jump Markov process  $(X_t, Y_t)_{t \in \mathbb{R}_+}$ . (b) Transition probabilities of the embedded random walk, which we denote by  $(X_t, Y_t)_{t \in \mathbb{N}}$  (here and throughout,  $\mathbb{N}$  is the set  $\{0, 1, 2, \dots\}$ ), with an abuse of notation.

$aa$ ) is  $A$  (resp.,  $A$  and  $a$ ). Only pollen and stigmata with different proteins can give viable seeds; i.e., pollen of a plant of phenotype  $A$  can only fecundate stigmata of a plant of phenotype  $a$  and vice versa. It can be seen that seeds  $AA$  cannot be created, since the genotype of individuals of phenotype  $a$  is necessarily  $aa$ , which combine only with individuals of phenotype  $A$  (that have genotypes  $AA$  or  $Aa$ ); therefore we can consider without restriction populations consisting only of individuals of genotypes  $Aa$  and  $aa$ . Each viable seed is then necessarily of genotype  $Aa$  or  $aa$  with probability  $1/2$ . It is assumed that ovules are produced in continuous time at rate  $r > 0$ , and that each ovule is fecundated to give a seed, provided that there exists compatible pollen in the population. The lifetime of each individual follows an exponential distribution with mean  $1/d$ , where  $d > 0$ . Throughout, we consider

$$(1.1) \quad r > d,$$

which, we will see, is the interesting case.

Let us denote by  $X_t$  and  $Y_t$  the numbers of individuals of genotype  $Aa$  (phenotype  $A$ ) and  $aa$  (phenotype  $a$ ) at time  $t \in \mathbb{R}_+$ . The process  $(X_t, Y_t)_{t \in \mathbb{R}_+}$  is a pure-jump Markov process with transitions represented in Figure 1.1(a). A stochastic differential equation (SDE) representation of  $(X_t, Y_t)_{t \in \mathbb{R}_+}$  is given in [4]. Here we forget the continuous-time process and are interested in the embedded discrete-time Markov chain, which we denote, with an abuse of notation, by  $(X_t, Y_t)_{t \in \mathbb{N}}$ , and with transitions represented in Figure 1.1(b):

$$\begin{aligned} \mathbb{P}_{i,j}[(X_1, Y_1) = (i - 1, j)] &= \frac{di}{(r + d)(i + j)}, & \mathbb{P}_{i,j}[(X_1, Y_1) = (i + 1, j)] &= \frac{r}{2(r + d)}, \\ \mathbb{P}_{i,j}[(X_1, Y_1) = (i, j - 1)] &= \frac{dj}{(r + d)(i + j)}, & \mathbb{P}_{i,j}[(X_1, Y_1) = (i, j + 1)] &= \frac{r}{2(r + d)}, \end{aligned}$$

where  $\mathbb{P}_{i,j}$  means that the process starts with the initial condition  $(X_0, Y_0) = (i, j)$ . The main and profound difficulty is that this random walk is not homogeneous in space, while techniques developed in the literature for random walks on positive quadrants mostly focus on the homogeneous case (see, e.g., Fayolle, Iasnogorodski, and Malyshev [6], Klein Haneveld and Pittenger [8], Kurkova and Raschel [9], and Walraevens, van Leeuwen, and Boxma [13]). We introduce a generating function

(1.4) that satisfies a PDE of a new type that we solve. Although the particularity of the problem is exploited, these techniques and the links between probability and PDEs may be extended to carry out general studies of inhomogeneous random walks in cones. The introduction of PDEs through generating functions had been already used by Feller [7] for a trunking problem with an inhomogeneous random walk in dimension 1. To our knowledge, the case of inhomogeneous random walks in the cone with absorbing boundaries has been left open. In [10], the discriminatory processor-sharing queue is considered, but boundaries are not absorbing and the overall arrival rate is constant, which is not the case in our model.

When one of the phenotypes  $A$  or  $a$  disappears, reproduction becomes impossible, and the system becomes extinct. We are interested in the probability of extinction of  $(X_t, Y_t)_{t \in \mathbb{N}}$  (or, equivalently, in that of  $(X_t, Y_t)_{t \in \mathbb{R}_+}$ ). Let us introduce the first time at which one of the two types becomes extinct:

$$(1.2) \quad \tau_0 = \inf\{t \in \mathbb{N} : X_t = 0 \text{ or } Y_t = 0\}.$$

For  $i, j \in \mathbb{N}$ , let us denote by

$$(1.3) \quad p_{i,j} = \mathbb{P}_{i,j}[\tau_0 < \infty]$$

the absorption probabilities, and by

$$(1.4) \quad P(x, y) = \sum_{i,j \geq 1} p_{i,j} x^i y^j$$

their generating function. By symmetry arguments, we have, for all  $i, j \in \mathbb{N}$ ,

$$(1.5) \quad p_{i,j} = p_{j,i}.$$

Moreover, for any  $i, j \in \mathbb{N}$  such that  $i = 0$  or  $j = 0$ , we have

$$(1.6) \quad p_{i,j} = 1.$$

In section 2, we will see that the  $p_{i,j}$ 's satisfy the Dirichlet problem associated with the doubly indexed recurrence equation

$$(1.7) \quad q_{i,j} = \frac{di}{(r+d)(i+j)} q_{i-1,j} + \frac{dj}{(r+d)(i+j)} q_{i,j-1} + \frac{r}{2(r+d)} q_{i,j+1} + \frac{r}{2(r+d)} q_{i+1,j}$$

and with the boundary condition (1.6). This problem does not admit simple solutions. There is no uniqueness of solutions to this problem. Note that the constant sequence equal to 1 is a solution. However, we are interested in solutions that tend to 0 as  $i$  or  $j$  tends to infinity, since (see [4] and Proposition 2.2 in this paper) estimates for  $p_{i,j}$  were obtained through probabilistic coupling techniques; they show that in the case (1.1) that we consider,  $p_{i,j}$  is strictly less than 1. In fact, the  $p_{i,j}$ 's correspond to the smallest positive solution of the Dirichlet problem and are completely determined if we give the probabilities  $(p_{i,1})_{i \geq 1}$ . We conclude the section with more precise estimates of the absorption probabilities  $p_{i,j}$  as the initial state  $(i, j)$  goes to infinity along one axis (Proposition 2.3). These new estimates rely on Proposition 2.2 and on comparisons with one-dimensional random walks. In section 3, we consider the generating function  $P(x, y)$  associated with the  $p_{i,j}$ 's and show that it satisfies a PDE, which has one and only one solution, that is computed (Proposition 3.8) explicitly with a dependence

on the  $(p_{i,1})_{i \geq 1}$ , prompting us to use the name “Green’s function.” This provides a new formulation of the solution of (1.7), that is, however, difficult to work with numerically. Hence, in section 4, we propose two different approaches leading to numerical approximations of the solution of the Dirichlet problem (1.6)–(1.7), which are based on stochastic and deterministic approaches.

In summary, we provide here several approaches to handling the extinction probabilities of the inhomogeneous random walk  $(X_t, Y_t)_{t \in \mathbb{R}_+}$  of our problem. Estimates from [4] are recalled, and the recurrence equation (1.7) is solved numerically and theoretically, pending further investigation of the PDE formulation.

**2. Existence of a solution.**

**2.1. Dirichlet problem.** We first establish that the extinction probabilities  $p_{i,j}$  of (1.3) solve the Dirichlet problem (1.6)–(1.7).

PROPOSITION 2.1.

- (i) *The extinction probabilities  $(p_{i,j})_{i,j \geq 1}$  are solutions to the Dirichlet problem (1.7) with boundary condition (1.6). Uniqueness of the solution may not hold, but the extinction probabilities  $(p_{i,j})_{i,j \in \mathbb{N}}$  define the smallest positive solution to this problem.*
- (ii) *Let the probabilities  $(p_{i,1})_{i \geq 1}$  be given. Then the probabilities  $(p_{i,j})_{i,j \geq 1}$  are completely determined.*

*Proof.* We begin with point (i). Equation (1.7) is obtained by using the strong Markov property at the time of the first event. Let us denote by  $K$  the transition kernel of the discrete-time Markov chain  $(X_t, Y_t)_{t \in \mathbb{N}^*}$ ; we have

$$Kf(i, j) = (f(i + 1, j) + f(i, j + 1)) \frac{r}{2(r + d)} + f(i - 1, j) \frac{di}{(r + d)(i + j)} + f(i, j - 1) \frac{dj}{(r + d)(i + j)}.$$

Following classical proofs (e.g., [3, 11]), the extinction probabilities  $(p_{i,j})_{i,j \in \mathbb{N}}$  satisfy the equation

$$(2.1) \quad \forall i, j \in \mathbb{N}^*, \quad f(i, j) = Kf(i, j) \quad \text{and} \quad \forall i, j \in \mathbb{N}, \quad f(i, 0) = f(0, j) = 1.$$

The constant solution equal to 1 is a solution to (2.1). Let us prove that  $(p_{i,j})_{i,j \in \mathbb{N}}$  is the smallest positive solution to (2.1). Let  $f$  be another positive solution. Let us consider  $M_t = f(X_{\inf\{t, \tau_0\}}, Y_{\inf\{t, \tau_0\}})$ , with  $\tau_0$  defined in (1.2). Denoting by  $(\mathcal{G}_t)_{t \in \mathbb{N}}$  the filtration of  $(M_t)_{t \in \mathbb{N}}$ , we have

$$\begin{aligned} \mathbb{E}[M_{t+1} | \mathcal{G}_t] &= \mathbb{E}[M_{t+1} \mathbb{1}_{\tau_0 \leq t} + M_{t+1} \mathbb{1}_{\tau_0 > t} | \mathcal{G}_t] \\ &= \mathbb{E}[M_t \mathbb{1}_{\tau_0 \leq t} + f(X_{t+1}, Y_{t+1}) \mathbb{1}_{\tau_0 > t} | \mathcal{G}_t] \\ &= M_t \mathbb{1}_{\tau_0 \leq t} + \mathbb{1}_{\tau_0 > t} \mathbb{E}[f(X_{t+1}, Y_{t+1}) | \mathcal{G}_t] \\ &= M_t \mathbb{1}_{\tau_0 \leq t} + \mathbb{1}_{\tau_0 > t} Kf(X_t, Y_t) \\ &= M_t \mathbb{1}_{\tau_0 \leq t} + \mathbb{1}_{\tau_0 > t} f(X_t, Y_t) = M_t. \end{aligned}$$

Hence  $(M_t)_{t \in \mathbb{N}}$  is a martingale, which converges on  $\{\tau_0 < \infty\}$  to  $f(X_{\tau_0}, Y_{\tau_0}) = 1$  (see the boundary condition in (2.1)). Thus by using the positivity of  $f$  and Fatou’s lemma, we obtain that for every  $i, j \in \mathbb{N}$

$$f(i, j) = \mathbb{E}_{i,j}[M_0] = \lim_{t \rightarrow \infty} \mathbb{E}_{i,j}[M_t] \geq \mathbb{E} \left[ \liminf_{t \rightarrow \infty} M_t \mathbb{1}_{\tau_0 < \infty} \right] = \mathbb{E}_{i,j}[\mathbb{1}_{\tau_0 < \infty}] = p_{i,j}.$$

This concludes the proof of (i).

Let us now consider point (ii). Assume that the probabilities  $(p_{i,1})_{i \geq 1}$  are given, and let us prove, by recursion, that every  $p_{i,j}$  can be computed. By symmetry, we need to prove only that this is the case for  $i \geq j$ . Assume

(Hrec  $j$ ): for  $j \in \mathbb{N}^*$  all the  $p_{k,\ell}$ 's for  $\ell \leq j$  and  $k \geq \ell$  can be computed from the  $p_{i,1}$ 's, and let us prove that we can determine the  $p_{i,j+1}$ 's for  $i \geq j + 1$ . From (1.7) we get

$$(2.2) \quad p_{i,j+1} = \frac{2(r+d)}{r} p_{i,j} - \frac{2di}{r(i+j)} p_{i-1,j} - \frac{2dj}{r(i+j)} p_{i,j-1} - p_{i+1,j}.$$

All the terms on the r.h.s. (right-hand side) of (2.2) are known by (Hrec  $j$ ), and hence  $p_{i,j+1}$  can be computed for any  $i \geq j + 1$ . This concludes the recursion.  $\square$

The following result shows that there is almost sure extinction in the case  $r \leq d$ . In the interesting case  $r > d$ , it also shows that there is a nontrivial solution to the Dirichlet problem (1.6)–(1.7).

PROPOSITION 2.2 (Proposition 9 of [4]). *We have the following regimes, given the parameters  $r$  and  $d$ :*

- (i) *If  $r \leq d$ , we have almost sure extinction of the population.*
- (ii) *If  $r > d (> 0)$ , then there is a strictly positive survival probability. Denoting by  $(i, j)$  the initial condition, we have*

$$\left(\frac{d}{r}\right)^{i+j} \leq p_{i,j} \leq \left(\frac{d}{r}\right)^i + \left(\frac{d}{r}\right)^j - \left(\frac{d}{r}\right)^{i+j}.$$

In point (ii), only bounds, and no explicit formulas, are available for the extinction probability  $p_{i,j}$ . The purpose of this article is to address (1.7) by considering the Green's function  $P(x, y)$  introduced in (1.4).

**2.2. Asymptotic behavior of the absorption probability as the initial state goes to infinity along one axis.** In this part, using the result of Proposition 2.2, we provide more precise estimates of the asymptotic behavior of the absorption probability  $p_{1,j} = p_{j,1}$  when  $j \rightarrow \infty$ . In particular, these estimates will be very useful when we tackle deterministic numerical simulations (see section 4.2).

PROPOSITION 2.3. *If  $j \rightarrow \infty$ , then*

$$(2.3) \quad p_{1,j} = p_{j,1} = \frac{2d}{r} \frac{1}{j} - \frac{2d(r^2 + dr + 2d^2)}{r^2(r+d)} \frac{1}{j^2} + O\left(\frac{1}{j^3}\right).$$

*Proof.* In addition to  $\tau_0$ , defined in (1.2), we introduce

$$S = \inf\{t \in \mathbb{N} : Y_t = 0\}, \quad T = \inf\{t \in \mathbb{N} : X_t = 0\},$$

the hitting times of the horizontal and vertical axes, respectively. Note that we have  $\tau_0 = \inf\{S, T\}$ . Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be a function such that  $f(j) < j$  for any  $j \geq 1$ . In what follows, we will choose  $f(j) = \lfloor \epsilon j \rfloor$ , with  $\epsilon \in (0, 1)$  and where  $\lfloor \cdot \rfloor$  denotes the integer part. We obviously have the identity

$$(2.4) \quad p_{1,j} = \mathbb{P}_{(1,j)}[\tau_0 < \infty] = \mathbb{P}_{(1,j)}[\tau_0 \leq f(j)] + \mathbb{P}_{(1,j)}[f(j) < \tau_0 < \infty].$$

To prove Proposition 2.3, we shall give estimates for both terms on the r.h.s. of (2.4).

*First step:* Study of  $\mathbb{P}_{(1,j)}[\tau_0 \leq f(j)]$ . Since  $f(j) < j$ , it is impossible, starting from  $(1, j)$ , to reach the horizontal axis before time  $f(j)$ , and we have  $\mathbb{P}_{(1,j)}[\tau_0 \leq f(j)] = \mathbb{P}_{(1,j)}[T \leq f(j)]$ . In order to compute the latter probability, we introduce two one-dimensional random walks on  $\mathbb{N}$ , namely  $X^-$  and  $X^+$ , which are killed at 0 and which have the jumps

$$\mathbb{P}_i[X_1^\pm = i - 1] = q_i^\pm, \quad \mathbb{P}_i[X_1^\pm = i + 1] = p_i^\pm, \quad \mathbb{P}_i[X_1^\pm = i] = r_i^\pm, \quad q_i^\pm + p_i^\pm + r_i^\pm = 1,$$

where

$$(2.5) \quad q_i^\pm = \frac{di}{(r+d)(i+j \mp f(j))}, \quad p_i^\pm = \frac{r}{2(r+d)}.$$

Both  $X^-$  and  $X^+$  are (inhomogeneous) birth-and-death processes on  $\mathbb{N}$ . These random walks are implicitly parameterized by  $j$ . If  $T^\pm = \inf\{t \in \mathbb{N} : X_t^\pm = 0\}$ , then

$$(2.6) \quad \mathbb{P}_1[T^- \leq f(j)] \leq \mathbb{P}_{(1,j)}[T \leq f(j)] \leq \mathbb{P}_1[T^+ \leq f(j)].$$

The quantities  $\mathbb{P}_1[T^\pm \leq f(j)]$  are computable: we shall prove that

$$(2.7) \quad \mathbb{P}_1[T^\pm \leq f(j)] = \frac{2d}{r} \frac{1}{(j \mp f(j))} - \frac{2d(r^2 + dr + 2d^2)}{r^2(r+d)} \frac{1}{(j \mp f(j))^2} + O\left(\frac{1}{(j \mp f(j))^3}\right).$$

The main idea for proving (2.7) is that, the  $q_i^\pm$  being very small as  $j \rightarrow \infty$ , the only paths which will significantly contribute to the probability  $\mathbb{P}_1[T^\pm \leq f(j)]$  are the ones with very few jumps to the right. Let us define

$$\begin{aligned} \Lambda_t^\pm(p) &= \{\text{the chain } X^\pm \text{ makes exactly } p \text{ jumps to the right between } 0 \text{ and } t\} \\ &= \{\text{there exist } 0 \leq q_1 < \dots < q_p \leq t - 1 \text{ such that} \\ &\quad X_{q_1+1}^\pm - X_{q_1}^\pm = \dots = X_{q_p+1}^\pm - X_{q_p}^\pm = 1\}. \end{aligned}$$

We are entitled to write

$$(2.8) \quad \mathbb{P}_1[T^\pm \leq f(j)] = \mathbb{P}_1[T^\pm \leq f(j), \Lambda_{f(j)}^\pm(0)] + \mathbb{P}_1[T^\pm \leq f(j), \Lambda_{f(j)}^\pm(1)] \\ + \mathbb{P}_1[T^\pm \leq f(j), \cup_{p \geq 2} \Lambda_{f(j)}^\pm(p)],$$

and we now separately analyze the three terms on the right-hand side of (2.8). First,

$$(2.9) \quad \mathbb{P}_1[T^\pm \leq f(j), \Lambda_{f(j)}^\pm(0)] = \sum_{k=1}^{f(j)} \mathbb{P}_1[T^\pm = k, \Lambda_{f(j)}^\pm(0)] = \sum_{k=1}^{f(j)} (r_1^\pm)^{k-1} q_1^\pm \\ = \frac{q_1^\pm}{1 - r_1^\pm} (1 - (r_1^\pm)^{f(j)}) = \frac{q_1^\pm}{1 - r_1^\pm} \left(1 + O\left((r_1^\pm)^{f(j)}\right)\right).$$

A Taylor expansion of  $q_1^\pm/(1 - r_1^\pm)$  according to the powers of  $1/(j \mp f(j))$  together with the fact that  $(r_1^\pm)^{f(j)} = o(1/(j \mp f(j))^3)$  provides that

$$(2.10) \quad \mathbb{P}_1[T^\pm \leq f(j), \Lambda_{f(j)}^\pm(0)] = \frac{2d}{r(j \mp f(j))} \left(1 - \frac{1 + \frac{2d}{r}}{j \mp f(j)} + O\left(\frac{1}{(j \mp f(j))^2}\right)\right).$$

We now consider the second term on the right-hand side of (2.8). For the event  $\Lambda_{f(j)}^\pm(1)$ ,  $X^\pm$  first stays a time  $k_1$  at 1, then jumps to 2, where it remains for  $k_2$  unit

of time; it next goes to 1, and, after a time  $k_3$ , jumps to 0. Further, since  $T^\pm \leq f(j)$ , we have  $k_1 + 1 + k_2 + 1 + k_3 + 1 \leq f(j)$ . Denoting by  $\tilde{k}_1 = k_1 + k_3$  the time spent in position 1, we thus have

$$\begin{aligned}
 (2.11) \quad & \mathbb{P}_1[T^\pm \leq f(j), \Lambda_{f(j)}^\pm(1)] \\
 &= \sum_{k_1+k_2+k_3 \leq f(j)-3} (r_1^\pm)^{k_1} p_1^\pm (r_2^\pm)^{k_2} q_2^\pm (r_1^\pm)^{k_3} q_1^\pm = p_1^\pm q_1^\pm q_2^\pm \sum_{\tilde{k}_1+k_2 \leq f(j)-3} (r_1^\pm)^{\tilde{k}_1} (r_2^\pm)^{k_2} \\
 &= \frac{p_1^\pm q_1^\pm q_2^\pm}{(1-r_1^\pm)(1-r_2^\pm)} \left(1 + O\left((r_1^\pm \vee r_2^\pm)^{f(j)-2}\right)\right).
 \end{aligned}$$

As for the first term, using the fact that  $(r_1^\pm \vee r_2^\pm)^{f(j)-2} = o(1/(j \mp f(j))^3)$  and a Taylor expansion according to the powers of  $1/(j \mp f(j))$  gives that

$$(2.12) \quad \mathbb{P}_1[T^\pm \leq f(j), \Lambda_{f(j)}^\pm(1)] = \frac{4d^2}{r(r+d)(j \mp f(j))^2} \left(1 + O\left(\frac{1}{(j \mp f(j))}\right)\right).$$

Finally, let us consider the third term  $\mathbb{P}_1[T^\pm \leq f(j), \cup_{p \geq 2} \Lambda_{f(j)}^\pm(p)]$ . On  $\cup_{p \geq 2} \Lambda_{f(j)}^\pm(p)$ , the two first jumps to the right are either from 1 to 2 and 2 to 3, or twice from 1 to 2. Thus, extinction means that there are at least three jumps from 3 to 2, 2 to 1, and 1 to 0 or two jumps from 2 to 1 and 1 to 0. Since  $q_i^\pm$  is an increasing function of  $i$ , we deduce

$$\begin{aligned}
 (2.13) \quad & \mathbb{P}_1[T^\pm \leq f(j), \cup_{p \geq 2} \Lambda_{f(j)}^\pm(p)] \\
 & \leq (q_3^\pm)^3 = \left(\frac{d}{r+d}\right)^3 \left(\frac{3}{3+j \mp f(j)}\right)^3 = O\left(\frac{1}{(j \mp f(j))^3}\right).
 \end{aligned}$$

From (2.8), (2.9), (2.11), and (2.13), we obtain (2.7).

*Second step:* Study of  $\mathbb{P}_{(1,j)}[f(j) < \tau_0 < \infty]$ . We have that

$$\begin{aligned}
 (2.14) \quad & \mathbb{P}_{(1,j)}[f(j) < \tau_0 < \infty] \\
 &= \sum_{k, \ell \geq 1} \mathbb{P}_{(1,j)}[f(j) < \tau_0 < \infty | (X_{f(j)}, Y_{f(j)}) = (k, \ell)] \mathbb{P}_{(1,j)}[(X_{f(j)}, Y_{f(j)}) = (k, \ell)] \\
 &= \sum_{k, \ell \geq 1} p_{k, \ell} \mathbb{P}_{(1,j)}[(X_{f(j)}, Y_{f(j)}) = (k, \ell)],
 \end{aligned}$$

by using the strong Markov property. Introduce now a function  $g : \mathbb{N} \rightarrow \mathbb{N}$  such that  $f(j) + g(j) < j$  for any  $j \geq 1$ , we can split (2.14) into

$$(2.15) \quad \sum_{k, \ell \geq g(j)} p_{k, \ell} \mathbb{P}_{(1,j)}[(X_{f(j)}, Y_{f(j)}) = (k, \ell)] + \sum_{\substack{g(j) \geq k \geq 1 \\ \text{and/or} \\ g(j) \geq \ell \geq 1}} p_{k, \ell} \mathbb{P}_{(1,j)}[(X_{f(j)}, Y_{f(j)}) = (k, \ell)].$$

With Proposition 2.2 we obtain the following upper bound for the first sum in (2.15):

$$\sum_{k, \ell \geq g(j)} p_{k, \ell} \mathbb{P}_{(1,j)}[(X_{f(j)}, Y_{f(j)}) = (k, \ell)] \leq 2 \left(\frac{d}{r}\right)^{g(j)}.$$

In particular, if we choose  $g$  such that as  $j \rightarrow \infty$ ,  $g(j) \rightarrow \infty$  fast enough, then clearly the term above is negligible compared to (2.7). For the second sum in (2.15),

$$\sum_{\substack{g(j) \geq k \geq 1 \\ \text{and/or} \\ g(j) \geq \ell \geq 1}} p_{k,\ell} \mathbb{P}_{(1,j)}[(X_{f(j)}, Y_{f(j)}) = (k, \ell)] = \sum_{g(j) \geq k \geq 1} p_{k,\ell} \mathbb{P}_{(1,j)}[(X_{f(j)}, Y_{f(j)}) = (k, \ell)],$$

since, by assumption,  $j - f(j) > g(j)$ , so that  $Y$  cannot reach values  $\ell \leq g(j)$  in  $f(j)$  steps. Then we have

$$(2.16) \quad \sum_{g(j) \geq k \geq 1} p_{k,\ell} \mathbb{P}_{(1,j)}[(X_{f(j)}, Y_{f(j)}) = (k, \ell)] \leq \sum_{g(j) \geq k \geq 1} \mathbb{P}_{(1,j)}[(X_{f(j)}, Y_{f(j)}) = (k, \ell)] \leq \mathbb{P}_{(1,j)}[0 \leq X_{f(j)} \leq g(j)].$$

To obtain an upper bound for (2.16) we are going to use, again, a one-dimensional random walk. Introduce  $\tilde{X}$ , a random walk on  $\mathbb{N}$ , which is killed at 0, homogeneous on  $\mathbb{N}^*$ , with jumps

$$\mathbb{P}_k[\tilde{X}_1 = k - 1] = \tilde{q}, \quad \mathbb{P}_k[\tilde{X}_1 = k + 1] = \tilde{p}, \quad \mathbb{P}_k[\tilde{X}_1 = k] = \tilde{r}, \quad \tilde{q} + \tilde{p} + \tilde{r} = 1,$$

where

$$\tilde{q} = \frac{d(1 + f(j))}{(r + d)(1 + j - 2f(j))}, \quad \tilde{p} = \frac{r}{2(r + d)}.$$

This walk is again parameterized by  $j$ . By construction of  $\tilde{X}$ , we have

$$\mathbb{P}_{(1,j)}[0 \leq X_{f(j)} \leq g(j)] \leq \mathbb{P}_1[\tilde{X}_{f(j)} \leq g(j)].$$

Denoting by  $\tilde{m}$  and  $\tilde{\sigma}^2$  the mean and variance of  $(\tilde{X}_2 - \tilde{X}_1)$ , respectively (they could easily be computed), we can write

$$\mathbb{P}_1[\tilde{X}_{f(j)} \leq g(j)] = \mathbb{P}_0 \left[ \frac{\tilde{X}_{f(j)} - \tilde{m}f(j)}{\tilde{\sigma}\sqrt{f(j)}} \leq \frac{g(j) - 1 - \tilde{m}f(j)}{\tilde{\sigma}\sqrt{f(j)}} \right].$$

By a suitable choice of the functions  $f$  and  $g$ , for instance  $f(j) = \lfloor \epsilon j \rfloor$  with  $\epsilon \in (0, 1)$  and  $g(j) = \lfloor j^{3/4} \rfloor$ , the central limit theorem gives that the latter is negligible compared to  $\mathbb{P}_{(1,j)}[\tau_0 \leq f(j)]$ . For this last term, using (2.7), (2.6) and letting  $\epsilon$  tend to 0 provides (2.3). The proof is concluded.  $\square$

Let us make some remarks on possible extensions of Proposition 2.3.

*Remark 2.4.* 1. The proof of Proposition 2.3 can easily be extended to the asymptotic of  $p_{i,j}$  as  $j \rightarrow \infty$  for any fixed value of  $i$ . In particular, we have the following asymptotic behavior:

$$(2.17) \quad p_{i,j} = \left(\frac{2d}{r}\right)^i \frac{i!}{j^i} + O\left(\frac{1}{j^{i+1}}\right).$$

2. It is possible to generalize (2.8) by

$$(2.18) \quad \mathbb{P}_1[T^\pm \leq f(j)] = \sum_{p=0}^{k-1} \mathbb{P}_1[T^\pm \leq f(j), \Lambda_p^{f(j)}] + \mathbb{P}_1[T^\pm \leq f(j), \cup_{p \geq k} \Lambda_p^{f(j)}].$$



We can show, as in the proof of Proposition 2.3, that  $\mathbb{P}_1[T^\pm \leq f(j), \cup_{p \geq k} \Lambda_p^{f(j)}] \leq (q_{k+1}^\pm)^{k+1} = o(1/j^{k+1})$  (see (2.13)) and that the probabilities  $\mathbb{P}_1[T^\pm \leq f(j), \Lambda_p^{f(j)}]$  admit Taylor expansions in powers of  $1/(j \mp f(j))$ , where the development for the  $p$ th probability has a main term in  $1/(j \mp f(j))^{p+1}$ . This allows us to push the developments in (2.7) to higher orders.

For instance, the next term in (2.3) can be obtained by a long computation. First, we generalize (2.11) by writing that

$$(2.19) \quad \mathbb{P}_1[T^\pm \leq f(j), \Lambda_2^{f(j)}] = q_1^\pm ((p_1^\pm)^2 (q_2^\pm)^2 + p_1^\pm p_2^\pm q_3^\pm q_2^\pm) \sum_{\tilde{k}_1 + \tilde{k}_2 + \tilde{k}_3 \leq f(j) - 5} (r_1^\pm)^{\tilde{k}_1} (r_2^\pm)^{\tilde{k}_2} (r_3^\pm)^{\tilde{k}_3},$$

where  $\tilde{k}_1, \tilde{k}_2$ , and  $\tilde{k}_3$  are the times spent by the random walk in the states 1, 2, and 3. Then, pushing the Taylor expansion further leads to

$$p_{1,j} = \frac{2d}{r} \frac{1}{j \mp f(j)} - \frac{2d(r^2 + dr + 2d^2)}{r^2(r+d)} \frac{1}{(j \mp f(j))^2} + d \left( \frac{2}{r} \left( 1 + \frac{2d}{r} \right)^2 - \frac{24d(\frac{r}{2} + d)}{r^2(r+d)} + \frac{5r^2 d^2}{2(r+d)^2(\frac{r}{2} + d)^3} \right) \frac{1}{(j \mp f(j))^3} + O \left( \frac{1}{(j \mp f(j))^4} \right).$$

### 3. Green’s function.

**3.1. A functional equation for the Green’s function.** In this section, we consider the Green’s function  $P(x, y)$  defined in (1.4), associated with a solution of (1.7) in the same spirit as what can be found in Feller [7, Chap. XVII]. We show that it satisfies a nonclassical linear PDE that can be solved (see Proposition 3.8).

PROPOSITION 3.1. (i) *The function  $P(x, y)$  formally satisfies*

$$(3.1) \quad AP(x, y) = h(x, y, P),$$

where

$$(3.2) \quad AP(x, y) = Q(x, y) \frac{\partial P}{\partial x}(x, y) + Q(y, x) \frac{\partial P}{\partial y}(x, y) + R(x, y)P(x, y),$$

$$Q(x, y) = (r + d)x - \frac{r}{2} - \frac{r}{2} \frac{x}{y} - dx^2,$$

$$R(x, y) = \frac{r}{2x} + \frac{r}{2y} - dx - dy,$$

and where

$$(3.3) \quad h(x, y, P) = -\frac{r}{2} \left( x \frac{\partial^2 P}{\partial x \partial y}(x, 0) + y \frac{\partial^2 P}{\partial y \partial x}(0, y) \right) + dxy \left( \frac{1}{1-x} + \frac{1}{1-y} \right).$$

(ii) *For given  $(p_{i,1})_{i \geq 1}$ , we have a unique classical solution to (3.1)–(3.3) on  $]0, 1[ \times ]0, 1[$ .*

The function  $h$  in (3.3) depends only on a boundary condition ( $\partial^2 P / \partial x \partial y$  at the boundaries  $x = 0$  or  $y = 0$ , i.e., the  $p_{i,1}$ ’s for  $i \in \mathbb{N}^*$ ), which is nonclassical, while the operator  $A$  is of first order and hence associated with some transport equations.

*Proof.* Let us first establish (i). Using the Markov property at time  $t = 1$ ,

$$p_{i,j} = \frac{r}{2(r+d)}(p_{i+1,j} + p_{i,j+1}) + \frac{dj}{(r+d)(i+j)}p_{i,j-1} + \frac{di}{(r+d)(i+j)}p_{i-1,j},$$

then multiplying by  $x^i y^j$ , and summing over  $i, j \in \mathbb{N}^*$  leads to

$$(3.4) \quad (r+d) \sum_{i,j \geq 1} (i+j)p_{i,j}x^i y^j = \frac{r}{2} \sum_{i,j \geq 1} (i+j)(p_{i+1,j} + p_{i,j+1})x^i y^j + d \sum_{i,j \geq 1} j p_{i,j-1}x^i y^j + d \sum_{i,j \geq 1} i p_{i-1,j}x^i y^j.$$

The l.h.s. of (3.4) equals

$$(3.5) \quad (r+d) \sum_{i,j \geq 1} (i+j)p_{i,j}x^i y^j = (r+d) \left( x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} \right) P(x, y).$$

For the r.h.s. of (3.4),

$$(3.6) \quad \begin{aligned} & \frac{r}{2} \sum_{\substack{j \geq 1 \\ i \geq 2}} p_{i,j}(i-1+j)x^{i-1}y^j + \frac{r}{2} \sum_{\substack{i \geq 1 \\ j \geq 2}} p_{i,j}(i+j-1)x^i y^{j-1} \\ & + d \sum_{\substack{i \geq 1 \\ j \geq 0}} (j+1)p_{i,j}x^i y^{j+1} + d \sum_{\substack{i \geq 0 \\ j \geq 1}} (i+1)p_{i,j}x^{i+1}y^j \\ & = \frac{r}{2} \sum_{\substack{j \geq 1 \\ i \geq 2}} p_{i,j}(i-1)x^{i-1}y^j + \frac{r}{2} \sum_{\substack{j \geq 1 \\ i \geq 2}} p_{i,j}jx^{i-1}y^j \\ & + \frac{r}{2} \sum_{\substack{i \geq 1 \\ j \geq 2}} p_{i,j}ix^i y^{j-1} + \frac{r}{2} \sum_{\substack{i \geq 1 \\ j \geq 2}} p_{i,j}(j-1)x^i y^{j-1} \\ & + d \sum_{\substack{i \geq 1 \\ j \geq 0}} (j+1)p_{i,j}x^i y^{j+1} + d \sum_{\substack{i \geq 0 \\ j \geq 1}} (i+1)p_{i,j}x^{i+1}y^j. \end{aligned}$$

For the first term on the r.h.s. of (3.6),

$$(3.7) \quad \begin{aligned} \frac{r}{2} \sum_{\substack{j \geq 1 \\ i \geq 2}} p_{i,j}(i-1)x^{i-1}y^j &= \frac{r}{2} \sum_{\substack{j \geq 1 \\ i \geq 1}} p_{i,j}(i-1)x^{i-1}y^j = \frac{r}{2} \sum_{\substack{j \geq 1 \\ i \geq 1}} p_{i,j}ix^{i-1}y^j - \frac{r}{2} \sum_{\substack{j \geq 1 \\ i \geq 1}} p_{i,j}x^{i-1}y^j \\ &= \frac{r}{2} \frac{\partial}{\partial x} P(x, y) - \frac{r}{2} \frac{P(x, y)}{x}. \end{aligned}$$

Similar computation holds for the fourth term of the r.h.s. of (3.6). For the second term,

$$(3.8) \quad \begin{aligned} \frac{r}{2} \sum_{\substack{j \geq 1 \\ i \geq 2}} p_{i,j}jx^{i-1}y^j &= \frac{r}{2} \frac{y}{x} \sum_{\substack{j \geq 1 \\ i \geq 2}} p_{i,j}jx^i y^{j-1} = \frac{r}{2} \frac{y}{x} \left( \sum_{\substack{j \geq 1 \\ i \geq 1}} p_{i,j}jx^i y^{j-1} - \sum_{j \geq 1} p_{1,j}jxy^{j-1} \right) \\ &= \frac{r}{2} \frac{y}{x} \frac{\partial}{\partial y} P(x, y) - \frac{r}{2} \sum_{j \geq 1} p_{1,j}jy^j. \end{aligned}$$

We handle the third term of the r.h.s. of (3.6) similarly. Now for the fifth term,

$$(3.9) \quad d \sum_{\substack{i \geq 1 \\ j \geq 0}} p_{i,j} (j+1) x^i y^{j+1} = d \left( y^2 \frac{\partial}{\partial y} P(x, y) + y P(x, y) + \sum_{i \geq 1} x^i y \right).$$

Similar computation holds for the last term of (3.6). From (3.5), (3.6), (3.7), (3.8), and (3.9) we deduce that

$$\begin{aligned} (r+d) \left( x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} \right) P(x, y) &= \frac{r}{2} \frac{\partial}{\partial x} P(x, y) - \frac{r}{2} \frac{P(x, y)}{x} + \frac{r}{2} \frac{\partial}{\partial y} P(x, y) - \frac{r}{2} \frac{P(x, y)}{y} \\ &+ \frac{r}{2} \frac{y}{x} \frac{\partial}{\partial y} P(x, y) - \frac{r}{2} \sum_{j \geq 1} p_{1,j} j y^j + \frac{r}{2} \frac{x}{y} \frac{\partial}{\partial x} P(x, y) - \frac{r}{2} \sum_{i \geq 1} p_{i,1} i x^i \\ &+ d \left( y^2 \frac{\partial}{\partial y} P(x, y) + y P(x, y) + \sum_{i \geq 1} x^i y \right) + d \left( x^2 \frac{\partial}{\partial x} P(x, y) + x P(x, y) + \sum_{j \geq 1} x y^j \right), \end{aligned}$$

and finally  $\sum_{i \geq 1} p_{i,1} i x^i = x \frac{\partial^2 P}{\partial x \partial y}(x, 0)$ .

For point (ii), local existence and uniqueness stem from classical theorems [14]. Note that we construct an explicit, albeit complicated, solution (3.8) using the method of characteristics. This concludes the proof.  $\square$

*Remark 3.2.* In the case of homogeneous random walks, the operator  $AP(x, y)$  has the product form  $R(x, y)P(x, y)$ ; see [9]. In some sense, this means that the inhomogeneity leads to partial derivatives in the functional equation.

*Remark 3.3.* This technique allowing us to compute the solution of a discrete linear problem thanks to generating series is also called the Z-transform method.

**3.2. Characteristic curves.** In sections 3.2 and 3.3 we establish, by the method of characteristic equations, an explicit formula for the solutions to (3.1), which proves the existence of the solution. Since  $A$  is a first-order differential operator, we have a transport-like PDE. We introduce the following characteristic ODEs. Let  $(x_s, y_s)_{s \in \mathbb{R}_+}$  be the solution to the system

$$(3.10) \quad \begin{cases} \dot{x}_s = \frac{dx}{ds}(s) = Q(x_s, y_s), \\ \dot{y}_s = \frac{dy}{ds}(s) = Q(y_s, x_s), \end{cases}$$

where  $Q$  has been defined in (3.2). The vector field  $(Q(x, y), Q(y, x))$  that drives the dynamical system (3.10) is represented in Figure 3.1. A neighborhood of points of interest is shown in Figure 3.2. The dynamical system (3.10) and its solutions will turn out to be decisive in what follows—e.g., in Proposition 3.7, where we will use these characteristic equations in order to express the solutions to the fundamental functional equation (3.1).

**PROPOSITION 3.4.** *For any initial condition  $(x_0, y_0) \in \mathbb{R}^2$  such that  $x_0, y_0 \neq 0$  there exists a unique solution to (3.10), defined for  $s \in \mathbb{R}$  by*

$$(3.11) \quad x_s = \frac{\lambda r e^{rs} + \mu d e^{ds}}{d(\lambda e^{rs} + \mu e^{ds} + 1)}, \quad y_s = \frac{\lambda r e^{rs} + \mu d e^{ds}}{d(\lambda e^{rs} + \mu e^{ds} - 1)},$$

with

$$(3.12) \quad \lambda = \frac{2d x_0 y_0 - d(x_0 + y_0)}{(x_0 - y_0)(r - d)}, \quad \mu = \frac{-2d x_0 y_0 + r(x_0 + y_0)}{(x_0 - y_0)(r - d)}.$$




FIG. 3.1. Differential system (3.10): Vector field  $(Q(x, y), Q(y, x))$ . The solid lines correspond to the sets  $\{Q(x, y) = 0\}$  or  $\{Q(y, x) = 0\}$ .




FIG. 3.2. Differential system (3.10). (a) Neighborhood of the saddle point  $(1, 1)$ . (b) Neighborhood of the attractive equilibrium  $(r/d, r/d)$ .

*Proof.* First, since  $Q(x, y)$  and  $Q(y, x)$  are locally Lipschitz-continuous for  $x, y \neq 0$ , we have local existence and uniqueness. Let us introduce the new variable  $z_s = x_s/y_s$ , which is defined as long as  $y_s \neq 0$ . Using the expression (3.2) of  $Q$ , the system (3.10) becomes

$$(3.13) \quad \dot{x}_s = (r + d)x_s - \frac{r}{2}(1 + z_s) - dx_s^2,$$

$$(3.14) \quad \begin{aligned} \dot{z}_s &= \frac{\dot{x}_s y_s - x_s \dot{y}_s}{y_s^2} \\ &= (r + d)z_s - \frac{r}{2} \frac{z_s}{x_s} - \frac{r}{2} \frac{z_s}{y_s} - dx_s z_s - (r + d)z_s + \frac{r}{2} \frac{z_s}{y_s} + \frac{r}{2} \frac{z_s}{x_s} + dy_s z_s \\ &= dx_s(1 - z_s). \end{aligned}$$

From (3.14), we obtain

$$(3.15) \quad x_s = \frac{\dot{z}_s}{d(1 - z_s)},$$

and differentiating (3.14) with respect to the time  $s$  gives

$$\dot{x}_s = \frac{d\dot{z}_s(1-z_s) + d\dot{z}_s^2}{d^2(1-z_s)^2}.$$

Plugging this expression and (3.15) into (3.13) provides

$$\frac{d\dot{z}_s(1-z_s) + d\dot{z}_s^2}{d^2(1-z_s)^2} = \frac{(r+d)\dot{z}_s}{d(1-z_s)} - \frac{r}{2}(1+z_s) - d\frac{\dot{z}_s^2}{d^2(1-z_s)^2},$$

and, therefore,

$$(3.16) \quad \ddot{z}_s(1-z_s) + 2\dot{z}_s^2 - (r+d)(1-z_s)\dot{z}_s + \frac{rd}{2}(1+z_s)(1-z_s)^2 = 0.$$

The system (3.15)–(3.16) can be solved explicitly. Let us define  $u_s = (1+z_s)/(1-z_s)$ , so that

$$(3.17) \quad z_s = \frac{u_s - 1}{u_s + 1},$$

from which we obtain

$$(3.18) \quad \dot{z}_s = \frac{2\dot{u}_s}{(1+u_s)^2}, \quad \ddot{z}_s = \frac{2\ddot{u}_s(1+u_s) - 4\dot{u}_s^2}{(1+u_s)^3}.$$

Using these expressions in (3.16) provides

$$(3.19) \quad \begin{aligned} & \frac{2\ddot{u}_s(1+u_s) - 4\dot{u}_s^2}{(1+u_s)^3} \frac{2}{u_s+1} + 2\frac{4\dot{u}_s^2}{(1+u_s)^4} \\ & - (r+d)\frac{2}{u_s+1} \frac{2\dot{u}_s}{(1+u_s)^2} + \frac{rd}{2} \frac{2u_s}{u_s+1} \frac{4}{(u_s+1)^2} = 0 \\ \Leftrightarrow & 4\ddot{u}_s(1+u_s) - 4(r+d)\dot{u}_s(1+u_s) + 4rd u_s(u_s+1) = 0 \\ \Leftrightarrow & \ddot{u}_s - (r+d)\dot{u}_s + rd u_s = 0. \end{aligned}$$

Hence  $u$  satisfies a second-order ODE, which admits the solutions

$$(3.20) \quad u_s = \lambda e^{rs} + \mu e^{ds}, \quad \lambda, \mu \in \mathbb{R}.$$

From (3.15), (3.17), and (3.20),

$$(3.21) \quad z_s = \frac{\lambda e^{rs} + \mu e^{ds} - 1}{\lambda e^{rs} + \mu e^{ds} + 1}, \quad x_s = \frac{\lambda r e^{rs} + \mu d e^{ds}}{d(\lambda e^{rs} + \mu e^{ds} + 1)}.$$

The integration constants  $\lambda$  and  $\mu$  can be expressed in terms of the initial conditions  $x_0$  and  $z_0 = x_0/y_0$ :

$$(3.22) \quad \lambda = \frac{2dx_0 - d(1+z_0)}{(1-z_0)(r-d)}, \quad \mu = \frac{r - 2dx_0 + z_0r}{(1-z_0)(r-d)}.$$

This yields the announced result with  $y_s = x_s/z_s$ .  $\square$




FIG. 3.3. Solutions corresponding to several initial conditions. We see that, depending on the initial condition, the solutions converge or diverge to infinity.

Remark 3.5. Explicit expressions for  $x_s$  and  $y_s$  as functions of time and parameterized by  $r$  and  $d$  can be obtained. Using the expressions (3.12) of  $\lambda$  and  $\mu$  in (3.21) and the relations between  $x_s$ ,  $y_s$ , and  $z_s$  provides

$$(3.23) \quad x_s = \frac{-\frac{x_0+y_0-2x_0y_0}{x_0+y_0-2(d/r)x_0y_0} \exp(rs) + \exp(ds)}{-\frac{d}{r} \frac{x_0+y_0-2x_0y_0}{x_0+y_0-2(d/r)x_0y_0} \exp(rs) + \exp(ds) + \frac{(1-d/r)(y_0-x_0)}{x_0+y_0-2(d/r)x_0y_0}},$$

$$(3.24) \quad y_s = \frac{-\frac{x_0+y_0-2x_0y_0}{x_0+y_0-2(d/r)x_0y_0} \exp(rs) + \exp(ds)}{-\frac{d}{r} \frac{x_0+y_0-2x_0y_0}{x_0+y_0-2(d/r)x_0y_0} \exp(rs) + \exp(ds) - \frac{(1-d/r)(y_0-x_0)}{x_0+y_0-2(d/r)x_0y_0}}.$$

For what follows, it is useful to notice that the constant

$$(3.25) \quad -\frac{\lambda r}{\mu d} = \frac{x_0 + y_0 - 2x_0y_0}{x_0 + y_0 - 2(d/r)x_0y_0},$$

which appears in (3.23) and (3.24), belongs to  $]0, 1[$  for all  $(x_0, y_0) \in ]0, 1[^2$ ; this is due to the fact that  $d/r < 1$ .

Below, we list some properties of the solutions to the dynamical system (3.10). The trajectories of the solutions to (3.10) can be decomposed into four steps. Several solutions, started with various initial conditions, are represented in Figure 3.3. In order to describe them, let us introduce

$$(3.26) \quad s_0 = \frac{1}{r-d} \log\left(-\frac{\mu d}{\lambda r}\right) = \frac{\log\left(\frac{x_0+y_0-2x_0y_0}{x_0+y_0-2(d/r)x_0y_0}\right)}{d-r}$$

and  $s_{\pm}$  as the only positive roots of the denominators of  $x_s$  and  $y_s$  in (3.11):

$$(3.27) \quad \lambda e^{rs} + \mu e^{ds} \pm 1.$$

PROPOSITION 3.6. Let  $(x_0, y_0) \in ]0, 1[^2$ .

- (i) The stationary solutions to (3.10) are the saddle point  $(1, 1)$  and the attractive point  $(r/d, r/d)$ .

- (ii)  $s_0 \in ]0, \infty[$  and  $s_{\pm} \in ]s_0, \infty[$ .
- (iii)  $\inf\{s_+, s_-\} = s_+$  (resp.,  $s_-$ ) if and only if  $y_0 < x_0$  (resp.,  $y_0 > x_0$ ).
- (iv)  $\lim_{s \uparrow s_+} x_s = -\infty$  and  $\lim_{s \downarrow s_-} y_s = -\infty$ .

The next points specify the behavior of the solution to (3.10).

- (v) On  $]0, s_0[$ ,  $(x_s, y_s)$  belongs to  $]0, 1[^2$  and goes to  $(x_{s_0}, y_{s_0}) = (0, 0)$  as  $s \rightarrow s_0$ .
- (vi) On  $]s_0, \inf\{s_+, s_-\}[$ ,  $(x_s, y_s)$  goes decreasingly to  $(x_{\inf\{s_+, s_-\}}, y_{\inf\{s_+, s_-\}})$ —by “decreasingly” we mean that both coordinates decrease.
- (vii) On  $] \inf\{s_+, s_-\}, \sup\{s_+, s_-\}[$ ,  $(x_s, y_s)$  goes to  $(x_{\sup\{s_+, s_-\}}, y_{\sup\{s_+, s_-\}})$  decreasingly.
- (viii) On  $] \sup\{s_+, s_-\}, \infty[$ ,  $(x_s, y_s)$  goes decreasingly to  $(r/d, r/d)$ .

*Proof of item (i).* To find the stationary solutions to (3.10), let us solve  $\dot{x} = 0$  and  $\dot{y} = 0$ . With (3.10), we get  $Q(x, y) = 0$  and  $Q(y, x) = 0$ . This directly implies that  $(x, y) = (1, 1)$  or  $(x, y) = (r/d, r/d)$ ; see Figures 3.1 and 3.2. Let us now study the stability of these two equilibria.

At  $(1, 1)$ , the Jacobian of (3.10) is

$$\text{Jac}(1, 1) = \frac{r}{2}J - dI,$$

where  $J$  is the  $2 \times 2$  matrix full of 1’s and where  $I$  is the  $2 \times 2$  identity matrix. The eigenvalues of  $\text{Jac}(1, 1)$  are  $-d < 0$  and  $r - d > 0$ , associated with the eigenvectors  $(1, -1)$  and  $(1, 1)$ , respectively. By classical linearization methods (e.g., [12, Chap. 3]), we deduce that the point  $(1, 1)$  is a saddle point.

At  $(r/d, r/d)$ , the Jacobian of (3.10) is

$$\text{Jac}(r/d, r/d) = \frac{d}{2}J - rI.$$

The eigenvalues are  $-r < 0$  and  $d - r < 0$ , associated with the eigenvectors  $(1, -1)$  and  $(1, 1)$ , respectively. The point  $(r/d, r/d)$  is therefore attractive.  $\square$

*Proof of items (ii)–(iv).* Now we prove the different facts dealing with  $s_0$ ,  $s_+$ , and  $s_-$ . First, (3.25) and the fact that  $r > d$  immediately imply that  $s_0 \in ]0, \infty[$ . Next, we show that (3.27) has on  $]0, \infty[$  only one root, which belongs to  $]s_0, \infty[$ . For this, we shall start with proving that (3.27) is positive on  $]0, s_0[$ . Then, we shall show that (3.27) is decreasing in  $]s_0, \infty[$  and goes to  $-\infty$  as  $s \rightarrow \infty$ .

In order to prove the first point above, it is enough to show that (3.27) is positive at  $s = 0$  and increasing on  $]0, s_0[$ . Expression (3.27) is positive at  $s = 0$  simply because

$$\begin{aligned} \lambda + \mu \pm 1 &= \frac{x_0 + y_0}{y_0 - x_0} \pm 1 = \frac{2y_0}{y_0 - x_0} \quad \text{or} \\ \frac{2x_0}{y_0 - x_0} &= \frac{(1 - d/r)[x_0 + y_0 \pm (y_0 - x_0)]}{x_0 + y_0 - 2(d/r)x_0y_0} > 0. \end{aligned}$$

To check that (3.27) is increasing on  $]0, s_0[$ , we note that the derivative of (3.27) is positive on  $]0, s_0[$ —actually by construction of  $s_0$ .

Now we prove the second point. From (3.25) and since  $r > d$ , we obtain that (3.27) goes to  $-\infty$  as  $s \rightarrow \infty$ . Also, by definition of  $s_0$ , the derivative of (3.27) is negative on  $]s_0, \infty[$ , and (3.27) is therefore decreasing on  $]s_0, \infty[$ .

The fact that  $\inf\{s_+, s_-\}$  equals  $s_+$  (resp.,  $s_-$ ) if and only if  $y_0 < x_0$  (resp.,  $y_0 > x_0$ ) follows directly from (3.27).

Finally, since the numerators of  $x_s$  and  $y_s$  are negative on  $]s_0, \infty[$ , hence in particular at  $s_{\pm}$ , it is immediate that  $\lim_{s \uparrow s_+} x_s = -\infty$  and  $\lim_{s \downarrow s_-} y_s = -\infty$ .  $\square$

*Proof of items (v)–(viii).* Let us first consider (v). By definition of  $s_0$ , the numerators of  $x_s$  and  $y_s$  in (3.23) and (3.24) vanish for the first time at  $s_0$ . Moreover, since  $s_{\pm} > s_0$ , both denominators are nonzero at  $s_0$  and  $x_{s_0} = y_{s_0} = 0$ . In particular, on  $]0, s_0[$ , we have  $(x_s, y_s) \in \mathbb{R}_+^2$ . In fact,  $(x_s, y_s) \in ]0, 1[^2$ . Indeed, on the segment  $\{1\} \times ]0, 1[$ ,  $\dot{x}_s < 0$ , whereas on  $]0, 1[ \times \{1\}$ ,  $\dot{y}_s < 0$ ; it is therefore not possible to go through these segments.

We turn to the proof of (vi). Thanks to (3.23) and (3.24), just after the time  $s_0$ ,  $(x_s, y_s)$  belongs to the negative quadrant  $\mathbb{R}_-^2$ . But for any  $(x, y) \in \mathbb{R}_-^2$ ,  $Q(x, y) \leq 0$  and  $Q(y, x) \leq 0$  (see Figure 3.1), in such a way that both  $x_s$  and  $y_s$  are decreasing as soon as they stay in this quarter plane, in other words for  $s \in ]s_0, \inf\{s_+, s_-\}[$ . At time  $\inf\{s_+, s_-\}$ , one (or even the two if  $s_+ = s_-$ , i.e., if  $x_0 = y_0$ ) of  $x_s$  and  $y_s$  becomes infinite. In the remainder, let us assume that  $\inf\{s_+, s_-\} = s_+$ ; similar reasoning would hold for the symmetrical case  $\inf\{s_+, s_-\} = s_-$ .

Let us show (vii). Just after  $s_+$ ,  $(x_s, y_s) \in (\mathbb{R}_+ \times \mathbb{R}_-) \cap \{(x, y) \in \mathbb{R}^2 : Q(x, y) < 0, Q(y, x) < 0\}$ . The latter set is simply connected and bounded by the curve  $\{(x, y) \in \mathbb{R}^2 : Q(x, y) = 0\}$ ; see Figure 3.1. Using classical arguments (see, e.g., [12]), we obtain that it is not possible to go through this limiting curve on which  $\dot{x}_s = 0$ ; this is why for any  $s \in ]s_+, s_-[$ ,  $(x_s, y_s)$  remains inside of this set.

We conclude with the proof of (viii). Just after the time  $s_-$ ,  $(x_s, y_s) \in \mathbb{R}_+^2 \cap \{(x, y) \in \mathbb{R}^2 : Q(x, y) < 0, Q(y, x) < 0\}$ . For the same reasons as above,  $(x_s, y_s)$  cannot leave this set and actually converges to  $(r/d, r/d)$ .  $\square$

**3.3. Use of the characteristic curves to simplify the functional equation.**

Let us assume the existence of a solution  $P(x, y)$  to (3.1), and let us define  $g_s = P(x_s, y_s)$ . Then

$$\begin{aligned} \dot{g}_s &= \frac{dg}{ds}(s) = \frac{\partial P}{\partial x}(x_s, y_s) \frac{dx_s}{ds} + \frac{\partial P}{\partial y}(x_s, y_s) \frac{dy_s}{ds} \\ &= \frac{\partial P}{\partial x}(x_s, y_s) Q(x_s, y_s) + \frac{\partial P}{\partial y}(x_s, y_s) Q(y_s, x_s). \end{aligned}$$

Thus, if  $P$  is a solution to (3.1), then

$$(3.28) \quad \dot{g}_s + R(x_s, y_s)g_s = h(x_s, y_s, P),$$

which looks like a first-order ODE for  $g$ , except that  $h$  depends on the boundary condition of  $P$ .

We first freeze the dependence on the solution in  $h$ ; i.e., we solve the ODE (3.28) as if the term on the r.h.s. were a known function. Using the solutions to the characteristic equations, we shall prove the following result.

**PROPOSITION 3.7.** *Let  $h(x, y)$  be an analytical function on  $]0, 1[^2$ . Let  $(x_0, y_0) \in \mathbb{R}^2$ . The solution to the ODE*

$$(3.29) \quad \dot{g}_s + R(x_s, y_s)g_s = h(x_s, y_s), \quad g_0 = P(x_0, y_0),$$

where  $(x_s, y_s)_{s \geq 0}$  are the solutions (3.11) to the characteristic curve starting at  $(x_0, y_0)$ , is given by

$$\begin{aligned} g_s^h &= P(x_0, y_0) \exp\left(-\int_0^s R(x_u, y_u) du\right) + \int_0^s h(x_u, y_u) e^{-\int_u^s R(x_\alpha, y_\alpha) d\alpha} du \\ (3.30) \quad &=: F(s, x_0, y_0, h). \end{aligned}$$



*Proof.* Equation (3.29) is an inhomogeneous first-order ODE. The solution to the associated homogeneous equation is

$$g_s = R(x_0, y_0) \exp\left(-\int_0^s R(x_u, y_u) du\right).$$

The announced result is deduced from the variation of constants method.  $\square$

A solution  $P$  to (3.1) hence satisfies the following functional equation for all  $s, x_0$ , and  $y_0$ :

$$(3.31) \quad P(x_s, y_s) = P(x_0, y_0) e^{-\int_0^s R(x_u, y_u) du} + \int_0^s h(x_u, y_u, P) e^{-\int_u^s R(x_\alpha, y_\alpha) d\alpha} du,$$

with the function  $h$  defined in (3.3). Plugging the definitions (1.4) and (3.3) into (3.31), we obtain

$$(3.32) \quad \sum_{i,j \geq 1} p_{i,j} x_s^i y_s^j = P(x_0, y_0) e^{-\int_0^s R(x_\alpha, y_\alpha) d\alpha} - \frac{r}{2} \sum_{i \geq 1} p_{i,1} \int_0^s i(x_u)^i e^{-\int_u^s R(x_\alpha, y_\alpha) d\alpha} du - \frac{r}{2} \sum_{j \geq 1} p_{1,j} \int_0^s j(y_u)^j e^{-\int_u^s R(x_\alpha, y_\alpha) d\alpha} du + d \int_0^s x_u y_u \left(\frac{1}{1-x_u} + \frac{1}{1-y_u}\right) e^{-\int_u^s R(x_\alpha, y_\alpha) d\alpha} du.$$

Notice that the r.h.s. of (3.32) depends only on the  $p_{i,1}$ 's and  $p_{1,j}$ 's, while the l.h.s. depends on all  $p_{i,j}$ 's.

PROPOSITION 3.8. *Let  $s_0 > 0$  be defined in (3.26). We have*

$$(3.33) \quad \begin{aligned} P(x_0, y_0) &= \frac{r}{2} \sum_{i \geq 1} p_{i,1} \int_0^{s_0} i(x_u)^i e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha} du \\ &\quad + \frac{r}{2} \sum_{j \geq 1} p_{1,j} \int_0^{s_0} j(y_u)^j e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha} du \\ &\quad - d \int_0^{s_0} x_u y_u \left(\frac{1}{1-x_u} + \frac{1}{1-y_u}\right) e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha} du. \end{aligned}$$

Before proving Proposition 3.8, let us show that the different quantities that appear in its statement are well defined. Indeed, this is a priori not clear: as  $\alpha \rightarrow s_0, x_\alpha \rightarrow 0$  and  $y_\alpha \rightarrow 0$ , in such a way that  $R(x_\alpha, y_\alpha) \rightarrow \infty$ ; see (3.2).

LEMMA 3.9. *Let  $i, j \in \mathbb{N}$ . Then  $\lim_{u \rightarrow s_0} (x_u)^i (y_u)^j e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha}$  is finite if and only if  $i + j \geq 1$ , and equals zero if and only if  $i + j \geq 2$ .*

*Proof.* First, since the only zero of any function of the form  $\alpha \exp(as) + \beta \exp(bs)$  with  $\alpha\beta < 0$  and  $a \neq b$  has order one, the following function has a simple zero at  $s_0$ :

$$(3.34) \quad -\frac{x_0 + y_0 - 2x_0y_0}{x_0 + y_0 - 2(d/r)x_0y_0} \exp(ru) + \exp(du).$$

Thanks to this and since  $s_+, s_- > s_0$ , both  $x_s$  and  $y_s$  have a zero of order one at  $s_0$ .

Moreover, with  $\lambda$  and  $\mu$  defined in (3.12), we obtain that

$$(3.35) \quad \begin{aligned} \exp\left(\int_0^u R(x_\alpha, y_\alpha) d\alpha\right) &= \frac{\lambda/\mu + 1 - 1/\mu}{(\lambda/\mu) \exp(ru) + \exp(du) - 1/\mu} \\ &\times \left(\frac{\lambda/\mu + 1 + 1/\mu}{(\lambda/\mu) \exp(ru) + \exp(du) + 1/\mu}\right) \frac{(r/d)(\lambda/\mu) + 1}{(r/d)(\lambda/\mu) \exp(ru) + \exp(du)} \exp((r+d)u). \end{aligned}$$

It is indeed easy to check that the derivative of the logarithm of (3.35) is equal to  $R(x_u, y_u)$ , for which we have an explicit expression; see (3.2), (3.23), and (3.24).

From (3.35), we see that  $e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha}$  has three poles, namely at  $s_0, s_+, s_-$ . The zero at  $s_0$  has order one by using again the considerations on the zeros of (3.34). In particular, Lemma 3.9 follows immediately.  $\square$

*Proof of Proposition 3.8.* Start by multiplying (3.32) by  $e^{\int_0^s R(x_\alpha, y_\alpha) d\alpha}$ , and then let  $s \rightarrow s_0$ . Since  $P(x, y) = xy \sum_{i,j \geq 1} p_{i,j} x^{i-1} y^{j-1}$  (see (1.4)), and since  $\lim_{s \rightarrow s_0} x_s y_s e^{\int_0^s R(x_\alpha, y_\alpha) d\alpha} = 0$  (see Lemma 3.9), we obtain that

$$\lim_{s \rightarrow s_0} P(x_s, y_s) e^{\int_0^s R(x_\alpha, y_\alpha) d\alpha} = 0,$$

which concludes the proof of Proposition 3.8.  $\square$

*Remark 3.10.* When  $(x_0, y_0) \in (0, 1)^2$ , we also have  $(x_u, y_u) \in (0, 1)^2$  for all  $u \in (0, s_0)$ . Thus it is possible to plug approximations of the  $p_{1,j}$ 's and  $p_{i,1}$ 's into (3.33) thanks to the terms  $(x_u)^i$  and  $(y_u)^j$ . Using that  $p_{1,i} = 2d/(ri) + o(1/i)$  when  $i \rightarrow +\infty$ , it is possible to find  $I_0$  sufficiently large so that

$$\begin{aligned} & \frac{r}{2} \sum_{i > I_0} p_{i,1} \int_0^{s_0} i((x_u)^i + (y_u)^i) e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha} du \\ & \sim d \int_0^{s_0} \sum_{i > I_0} ((x_u)^i + (y_u)^i) e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha} du \\ & = d \int_0^{s_0} \left( \frac{x_u^{I_0+1}}{1-x_u} + \frac{y_u^{I_0+1}}{1-y_u} \right) e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha} du. \end{aligned}$$

Thus

$$\begin{aligned} (3.36) \quad P(x_0, y_0) &= \frac{r}{2} \sum_{i=1}^{I_0} p_{i,1} \int_0^{s_0} i((x_u)^i + (y_u)^i) e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha} du \\ &+ d \int_0^{s_0} \left( \frac{x_u(x_u^{I_0} - y_u)}{1-x_u} + \frac{y_u(y_u^{I_0} - x_u)}{1-y_u} \right) e^{\int_0^u R(x_\alpha, y_\alpha) d\alpha} du + o(x_0^{I_0+1} + y_0^{I_0+1}). \end{aligned}$$

The latter expression shows that, numerically, one can restrict to the computation of a finite number of probabilities  $p_{i,1}$  for  $i \leq I_0$ .

**4. Numerical results.** In this section, we present two different ways of approximating the extinction probabilities  $p_{i,j}$ .

**4.1. Probabilistic algorithm.** A first possibility, if we are interested in a given initial condition  $(i, j)$ , is to approximate  $p_{i,j}$  by Monte Carlo simulations. For  $T > 0$  large, we simulate  $M$  paths  $(X_t^\ell, Y_t^\ell)_{t \in \{1, \dots, T\}}$  started at  $(i, j)$ , for  $\ell \in \{1, \dots, M\}$ , independent and distributed as the process  $(X_t, Y_t)_{t \in \{1, \dots, T\}}$ . The extinction probability is estimated by

$$\hat{p}_{M,T} = \frac{1}{M} \sum_{\ell=1}^M \mathbb{1}_{\{\exists t \leq T, X_t^\ell Y_t^\ell = 0\}}.$$

The estimator  $\hat{p}_{M,T}$  is the proportion of paths that have gone extinct before time  $T$ .

**PROPOSITION 4.1.** *Let  $(i, j)$  be the initial condition. The estimator  $\hat{p}_{M,T}$  has the following properties:*

- (i) It is a convergent and unbiased estimator of  $\mathbb{P}_{i,j}[\tau_0 \leq T]$ .
- (ii) Its variance is  $\mathbb{P}_{i,j}[\tau_0 \leq T](1 - \mathbb{P}_{i,j}[\tau_0 \leq T])/M$ , and hence we have the following asymptotic 95% confidence interval for  $p_{i,j}$ :

$$(4.1) \quad \left[ \widehat{p}_{M,T} - 1.96 \sqrt{\frac{\widehat{p}_{M,T}(1 - \widehat{p}_{M,T})}{M}}; \widehat{p}_{M,T} + 1.96 \sqrt{\frac{\widehat{p}_{M,T}(1 - \widehat{p}_{M,T})}{M}} \right].$$

*Proof.* These results are straightforward consequences of the law of large numbers and the central limit theorem, given that  $\mathbb{1}_{\{\exists t \leq T, X_t^\ell Y_t^\ell = 0\}}$  are independent Bernoulli random variables with parameter  $\mathbb{P}_{i,j}[\tau_0 \leq T]$ .  $\square$

Computing the extinction probabilities by Monte Carlo methods yields good results if we are interested in a given initial condition  $(i, j)$ . We then have a complexity of order  $M \times T$ . However, biologists may be interested in investigating the extinction probabilities when the initial condition  $(i, j)$  varies, and then that method become computationally expensive.

**4.2. Deterministic algorithm.** For numerical approximations, we restrict ourselves to the computation of  $(p_{i,j})_{i,j \in \{1, \dots, N\}}$  for a positive (large) integer  $N$ . In this case, (1.7) can be approximated by the solution to a linear system.

Let us define  $\mathbf{p}_N = (p_{1,1}, \dots, p_{1,N}, p_{2,1}, \dots, p_{2,N}, \dots, p_{N,1}, \dots, p_{N,N})^T$ , and let  $T_N$  be an  $N^2 \times N^2$  matrix with five nonzero diagonals:

$$T_N = \begin{pmatrix} A_1 & D & 0 & \dots & 0 \\ B_{2,1} & A_2 & D & \ddots & \vdots \\ 0 & B_{3,2} & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & D \\ 0 & \dots & 0 & B_{N,N-1} & A_N \end{pmatrix},$$

where  $D = \frac{r}{2(r+d)} Id_N$ ,  $A_i$  ( $i \in \{1, \dots, N\}$ ), and  $B_{i,i-1}$  ( $i \in \{2, \dots, N\}$ ) are the  $N \times N$  matrices

$$A_i = \begin{pmatrix} -1 & \frac{r}{2(r+d)} & 0 & \dots & 0 \\ \frac{d}{r+d} \frac{2}{i+2} & -1 & \ddots & \ddots & \vdots \\ 0 & \frac{d}{r+d} \frac{3}{i+3} & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & \frac{r}{2(r+d)} \\ 0 & \dots & 0 & \frac{d}{r+d} \frac{N}{i+N} & -1 \end{pmatrix},$$

$$B_{i,i-1} = \begin{pmatrix} \frac{d}{r+d} \frac{i}{i+1} & 0 & \dots & 0 \\ 0 & \frac{d}{r+d} \frac{i}{i+2} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & \frac{d}{r+d} \frac{i}{i+N} \end{pmatrix}.$$




FIG. 4.1. Estimation of the extinction probabilities  $p_{i,j}$  when  $r = 3$  and  $d = 2$ : (a) with the Monte Carlo method of section 4.1; (b) with the deterministic method of section 4.2.

Let us also define the vector  $b_N = (b_{1N}, \dots, b_{NN})^T \in \mathbb{R}^{N \times N}$  such that

$$\begin{aligned}
 b_{1N} &= - \begin{pmatrix} -\frac{d}{r+d} \\ \frac{d}{r+d} \frac{1}{1+2} \\ \vdots \\ \frac{d}{r+d} \frac{1}{1+(N-1)} \\ \frac{d}{r+d} \frac{1}{1+N} + \frac{r}{2(r+d)} \tilde{p}_{1,N+1} \end{pmatrix}, \\
 b_{iN} &= - \begin{pmatrix} \frac{d}{r+d} \frac{1}{1+i} \\ 0 \\ \vdots \\ 0 \\ \frac{r}{2(r+d)} \tilde{p}_{i,N+1} \end{pmatrix} \quad \text{for } i \in \{2, \dots, N-1\}, \\
 b_{NN} &= - \begin{pmatrix} \frac{d}{r+d} \frac{1}{1+N} + \frac{r}{2(r+d)} \tilde{p}_{N+1,1} \\ \frac{r}{2(r+d)} \tilde{p}_{N+1,2} \\ \vdots \\ \frac{r}{2(r+d)} \tilde{p}_{N+1,N-1} \\ \frac{r}{2(r+d)} (\tilde{p}_{N+1,N} + \tilde{p}_{N,N+1}) \end{pmatrix},
 \end{aligned}$$

where  $\tilde{p}_{i,N+1}$  and  $\tilde{p}_{N+1,N}$  are approximations of  $p_{i,N+1}$  given by Proposition 2.3. With these definitions, (1.7) can be rewritten as

$$T_N \mathbf{p}_N = b_N.$$

**4.3. Results.** We start with  $r = 3$  and  $d = 2$ . For the Monte Carlo simulation, we use  $M = 200$  and  $T = 5000$ . For the deterministic method, we use  $N = 50$ , so that  $(i, j) \in \{1, \dots, 50\}^2$ . Estimators of the extinction probabilities  $\hat{p}_{i,j}^{(1)}$  and  $\hat{p}_{i,j}^{(2)}$  obtained respectively from the methods of sections 4.1 and 4.2 are plotted in Figure 4.1. The results given by both methods are very similar, as shown by the statistics of Table 4.1. In Table 4.1, we compute the square difference between the two predictions  $(\hat{p}_{i,j}^{(1)} - \hat{p}_{i,j}^{(2)})^2$ , the absolute difference  $|\hat{p}_{i,j}^{(1)} - \hat{p}_{i,j}^{(2)}|$ , and the relative difference  $|\hat{p}_{i,j}^{(1)} - \hat{p}_{i,j}^{(2)}|/\hat{p}_{i,j}^{(2)}$ .

TABLE 4.1

Square, absolute, and relative differences between the predictions of the stochastic method when  $r = 3$  and  $d = 2$  (section 4.1) and of the deterministic methods (section 4.2). Recall that with  $M = 200$ , the width of the confidence interval (4.1) is  $6.92 \cdot 10^{-2}$ .

	Mean	St.dev.	Min	Max
Square error	$3.24 \cdot 10^{-5}$	$2.47 \cdot 10^{-4}$	$1.68 \cdot 10^{-36}$	$4.63 \cdot 10^{-3}$
Absolute error	$1.34 \cdot 10^{-3}$	$5.53 \cdot 10^{-3}$	$1.30 \cdot 10^{-18}$	$6.81 \cdot 10^{-2}$
Relative error	$4.49 \cdot 10^{-2}$	$1.71 \cdot 10^{-1}$	$9.80 \cdot 10^{-3}$	$9.71 \cdot 10^{-1}$




FIG. 4.2. Estimation of the extinction probabilities  $p_{i,j}$  when  $r = 2.002$  and  $d = 2$ : (a) with the Monte Carlo method of section 4.1; (b) with the deterministic method of section 4.2.

For the last of these, we consider only the couples  $(i, j)$ , where  $\hat{p}_{i,j}^{(1)}$  and  $\hat{p}_{i,j}^{(2)}$  do not vanish (else, the fraction is either not defined or is 1 whatever the value of  $\hat{p}_{i,j}^{(1)}$ ).

To carry further the comparison of the stochastic and deterministic methods, and to observe the influence of  $N$  on the quality of the approximation, we compute the relative quadratic error

$$(4.2) \quad \frac{\sqrt{\sum_{1 \leq i,j \leq 10} (\hat{p}_{i,j}^{(2)} - \hat{p}_{i,j}^{(3)})^2}}{\sqrt{\sum_{1 \leq i,j \leq 10} (\hat{p}_{i,j}^{(2)})^2}} \quad \text{or} \quad \frac{\sqrt{\sum_{1 \leq i,j \leq 10} (\hat{p}_{i,j}^{(2)} - \hat{p}_{i,j}^{(1)})^2}}{\sqrt{\sum_{1 \leq i,j \leq 10} (\hat{p}_{i,j}^{(3)})^2}}$$

when  $\hat{p}_{i,j}^{(2)}$  is the deterministic approximation for  $N \in \{10, \dots, 50\}$  and  $\hat{p}_{i,j}^{(3)}$  is given either by the deterministic approximation with  $N = 50$  or by the stochastic approximation  $\hat{p}_{i,j}^{(1)}$  with  $M = 200$ . In the first case, when  $\hat{p}_{i,j}^{(3)} = \hat{p}_{i,j}^{(1)}$ , the decrease in the quadratic errors stops around  $N = 18$  around 0.0891. This corresponds roughly to the stochastic error of the law of large numbers (4.1), which depends only on  $M$ . In the second case, when  $\hat{p}_{i,j}^{(3)}$  is the deterministic approximation with  $N = 50$ , the relative quadratic errors decrease exponentially fast in  $\exp(-0.6842 N)$  ( $R^2 = 99.92\%$ ).

In a second experiment, we choose  $r = 2.002$  and  $d = 2$ . This case is more interesting in population ecology, since small populations are of interest when they are fragile and endangered species. For the Monte Carlo simulation, we use  $M = 200$  and  $T = 5000$ . For the deterministic method, we use  $N = 100$ , so that  $(i, j) \in \{1, \dots, 100\}^2$ . The estimated extinction probabilities  $\hat{p}_{i,j}^{(1)}$  and  $\hat{p}_{i,j}^{(2)}$  are plotted in Figure 4.2, and statistics are computed in Table 4.2. Again, results from both methods are similar. This is confirmed by computing the relative quadratic errors, with the  $\hat{p}_{i,j}^{(2)}$ 's obtained

TABLE 4.2

Square, absolute, and relative differences between the predictions of the stochastic method when  $r = 2.002$  and  $d = 2$  (section 4.1) and of the deterministic methods (section 4.2). As in Table 4.1, since  $M = 200$ , the width of the confidence interval (4.1) is  $6.92 \cdot 10^{-2}$ .

	Mean	St.dev.	Min	Max
Square error	$6.27 \cdot 10^{-3}$	$6.25 \cdot 10^{-3}$	$8.42 \cdot 10^{-10}$	$5.31 \cdot 10^{-2}$
Absolute error	$6.89 \cdot 10^{-2}$	$3.90 \cdot 10^{-2}$	$2.90 \cdot 10^{-5}$	$2.31 \cdot 10^{-1}$
Relative error	$2.22 \cdot 10^{-1}$	$2.17 \cdot 10^{-1}$	$1.02 \cdot 10^{-4}$	1




FIG. 4.3. Estimation of the extinction probabilities  $p_{i,j}$  when  $r = 2.002$  and  $d = 2$ : Decrease with  $N$  of the log of the relative quadratic errors (4.2) between the deterministic and Monte Carlo methods ( $M = 200$ ).

from the deterministic method and the  $\hat{p}_{i,j}^{(3)} = \hat{p}_{i,j}^{(1)}$ 's from the Monte Carlo method. The decrease of this error is exponential with  $N$  in  $\exp(-0.0619 N)$  ( $R^2 = 98.98\%$ ) as shown in Figure 4.3. It can be noticed that, in this case, the performance of the Monte Carlo method match better that of the deterministic algorithm. This is due to the fact that Monte Carlo methods fail to produce good estimates of small probabilities (see [5] and references therein).

When the probabilities  $\hat{p}_{i,j}^{(3)}$ 's are given by the deterministic method with  $N = 50$  in (4.2), we have as in the previous case ( $r = 3$ ) an exponential decrease of the relative quadratic error in  $\exp(-0.1092 N)$  ( $R^2 = 95.07\%$ ).

**Acknowledgments.** The authors warmly thank the referees for useful comments and suggestions. P.L. thanks the members of her former affiliation, Laboratoire Paul Painlevé at the Université des Sciences et Technologies de Lille (USTL), for fruitful discussions and for giving her a productive working environment.

## REFERENCES

- [1] S. ASPANDIAROV, R. IASNOGORODSKI, AND M. MENSNIKOV, *Passage-time moments for non-negative stochastic processes and an application to reflected random walks in a quadrant*, Ann. Probab., 24 (1996), pp. 932–960.
- [2] K. B. ATHREYA AND P. E. NEY, *Branching Processes*, Springer, New York, 1970.
- [3] P. BALDI, L. MAZLIAK, AND P. PRIURET, *Martingales and Markov Chains*, Chapman & Hall, London, 2002.
- [4] S. BILLIARD AND V. C. TRAN, *A general stochastic model for sporophytic self-incompatibility*, J. Math. Biol., 64 (2012), pp. 163–210.
- [5] S. CLÉMENÇON, M. DÁVILA FELIPE, L.-V. LOZADA-CHANG, AND V. C. TRAN, *On computer-intensive simulation and estimation methods for rare event analysis in epidemic models*, manuscript, 2012.
- [6] G. FAYOLLE, R. IASNOGORODSKI, AND V. MALYSHEV, *Random Walks in the Quarter-Plane*, Springer-Verlag, Berlin, 1999.

- [7] W. FELLER, *An Introduction to Probability Theory and Its Applications*, Vol. I, John Wiley and Sons, New York, 1957.
- [8] L. A. KLEIN HANEVELD AND A. O. PITTENGER, *Escape time for a random walk from an orthant*, *Stochastic Process. Appl.*, 35 (1990), pp. 1–9.
- [9] I. KURKOVA AND K. RASCHEL, *Random walks in  $\mathbb{Z}_+^2$  with non-zero drift absorbed at the axes*, *Bull. Soc. Math. France*, 139 (2011), pp. 341–387.
- [10] K. M. REGE AND B. SENGUPTA, *Queue-length distribution for the discriminatory processor-sharing queue*, *Oper. Res.*, 44 (1996), pp. 653–657.
- [11] D. REVUZ, *Markov Chains*, North-Holland Math. Library, North-Holland, Amsterdam, 1984.
- [12] F. VERHULST, *Nonlinear Differential Equations and Dynamical Systems*, Springer Universitext, Springer, New York, 2000.
- [13] J. WALRAEVENS, J. S. H. VAN LEEUWAARDEN, AND O. J. BOXMA, *Power series approximations for two-class generalized processor sharing systems*, *Queueing Syst.*, 66 (2010), pp. 107–130.
- [14] E. ZACHMANOGLU AND D. THOE, *Introduction to Partial Differential Equations with Applications*, Dover Publications, New York, 1986.